

**PROSPECTIVE TWO YEARS STUDY OF COMPLICATIONS OF  
MANUAL SMALL INCISION CATARACT SURGERY IN EYES  
WITH PSEUDOEXFOLIATION SYNDROME**

**By**

**DR. PUNIT KUMAR SINGH**

**Dissertation Submitted to the**

**SRI DEVARAJ URS ACADEMY OF HIGHER EDUCATION AND RESEARCH**

**KOLAR**



In partial fulfillment

Of the requirements for the degree of

**MASTER OF SURGERY**

**In**

**OPHTHALMOLOGY**

**Under the guidance of**

**DR K. KANTHAMANI,**

**M.B.B.S., M.S.,**



**DEPARTMENT OF OPHTHALMOLOGY  
SRI DEVARAJ URS MEDICAL COLLEGE**

**Tamaka, Kolar**

**APRIL - 2011**

**SRI DEVARAJ URS ACADEMY OF HIGHER EDUCATION AND  
RESEARCH**

**DECLARATION BY THE CANDIDATE**

I hereby declare that this dissertation entitled  
**PROSPECTIVE TWO YEARS STUDY OF COMPLICATIONS OF  
MANUAL SMALL INCISION CATARACT SURGERY IN EYES  
WITH PSEUDOEXFOLIATION SYNDROME**

is a bonafide and genuine research work carried out by me under the guidance of  
**DR. K. KANTHAMANI M.B.B.S., M.S**, Professor, Department of Ophthalmology,  
Sri Devaraj Urs Medical College , Tamaka, Kolar in partial for the award of M.S  
degree in Ophthalmology to be held in 2011. This dissertation has not been submitted  
in part or full to any other university or towards any other degree before this below  
mentioned date.

**Date:**

**Signature of the Candidate**

**Place: Kolar**

**CERTIFICATE BY THE GUIDE**

This is to certify that the dissertation entitled  
**“PROSPECTIVE TWO YEARS STUDY OF COMPLICATIONS OF  
MANUAL SMALL INCISION CATARACT SURGERY IN EYES  
WITH PSEUDOEXFOLIATION SYNDROME**

" is a bonafide research work done by **DR. PUNIT KUMR SINGH** in partial fulfillment of the requirement for the degree of **MASTER OF SURGERY IN OPHTHALMOLOGY** as per regulations of **Sri Devaraj Urs Academy of Higher Education And Research, Kolar**. I have great pleasure in forwarding this to the university.

**Date:**

**Signature of the Guide**

**Place: Kolar**

**DR.K. KANTHAMANI**

**M.B.B.S., M.S,**

**Professor,**

**Department of Ophthalmology,**

**Sri Devaraj Urs Medical College,**

**Tamaka, Kolar**

**ENDORSEMENT BY THE HEAD OF THE DEPARTMENT**

This is to certify that the dissertation entitled

**“PROSPECTIVE TWO YEARS STUDY OF COMPLICATIONS OF  
MANUAL SMALL INCISION CATARACT SURGERY IN EYES  
WITH PSEUDOEXFOLIATION SYNDROME”**

is a bonafide research work done by **DR. PUNIT KUMAR SINGH.** under the guidance of

**Dr.K.KANTHAMANI, M.B.B.S., M.S,** Professor, Department of Ophthalmology,

Sri Devaraj Urs Medical College, Tamaka, Kolar.

Date:

Signature of the HOD

Place: Kolar

**DR. Narendra P Datti,**

MBBS,MS,

Professor and head of the department,

Department of Ophthalmology,

Sri Devaraj Urs Medical College,

Tamaka, Kolar

SRI DEVARAJ URS ACADEMY OF HIGHER EDUCATION AND RESEARCH

TAMAKA , KOLAR , KARNATAKA

**ENDORSEMENT BY THE HOD,  
PRINCIPAL / HEAD OF THE INSTITUTION**

This is to certify that the dissertation entitled  
**“PROSPECTIVE TWO YEARS STUDY OF COMPLICATIONS OF  
MANUAL SMALL INCISION CATARACT SURGERY IN EYES  
WITH PSEUDOEXFOLIATION SYNDROME”**

is a bonafide research work done by **DR. PUNIT KUMAR SINGH.**

under the guidance of **Dr.K.KANTHAMANI**, M.B.B.S., M.S, Professor,  
Department of Ophthalmology, Sri Devaraj Urs Medical College, Tamaka, Kolar.

**DR. Narendra P Datti,**  
MBBS,MS,  
Professor and head of the department,  
Department of Ophthalmology,  
Sri Devaraj Urs Medical College,  
Tamaka, Kolar

**Dr. M.B.Sanikop**  
**Principal**  
Sri Devaraj Urs Medical College  
Tamaka, Kolar

Date:

Place: Kolar

Date:

Place: Kolar

**SRI DEVARAJ URS ACADEMY OF HIGHER EDUCATION AND RESEARCH**

**TAMAKA , KOLAR , KARNATAKA**

**ETHICS COMMITTEE CERTIFICATE**

This is to certify that the Ethics committee of Sri Devaraj Urs  
Medical College, Tamaka, Kolar has unanimously approved

**DR. PUNIT KUMAR SINGH.**

Post-Graduate student in the subject of

**OPHTHALMOLOGY at**

**Sri Devaraj Urs Medical College, Kolar**

to take up the Dissertation work entitled

**“PROSPECTIVE TWO YEARS STUDY OF COMPLICATIONS OF MANUAL  
SMALL INCISION CATARACT SURGERY IN EYES WITH  
PSEUDOEXFOLIATION SYNDROME”**

to be submitted to the

**SRI DEVARAJ URS ACADEMY OF HIGHER EDUCATION AND RESEARCH**

**TAMAKA , KOLAR , KARNATAKA,**

**Member Secretary**

Sri Devaraj Urs Medical College,

Kolar–563101

Date :

Place : Kolar

## **COPYRIGHT**

### **DECLARATION BY THE CANDIDATE**

I hereby declare that the Sri Devaraj Urs Academy of Higher Education And Research, Kolar shall have the rights to preserve, use and disseminate this dissertation in print or electronic format for academic / research purpose.

Date:

Signature of the Candidate

**DR. PUNIT KUMAR SINGH**

**© Sri Devaraj Urs Academy of Higher Education And Research, Kolar**

## ACKNOWLEDGMENT

It is with great reverence, deep sense of gratitude and respect that I would like to thank my teacher and guide, DR. K.KANTHAMANI, M.B.B.S., M.S, Professor, Department of Ophthalmology, Sri Devraj Urs Medical College Tamaka, Kolar for her guidance, encouragement, and valuable insights during the entire period of this study and post graduation course.

I would like to express my appreciation and gratitude to Dr. Narendra P Datti, Professor and HOD, Department of Ophthalmology, Sri Devraj Urs Medical College Tamaka, Kolar, for his encouragement and suggestions during the entire course of this study and postgraduation course.

I want to express my profound gratitude to Dr.D. Krishnamurthy, Dr.Manjula my Professors Sri Devraj Urs Medical College Tamaka, Kolar whose knowledge and experience have guided me throughout my post graduation course.

I would like to express my heartfelt thanks to all my Assistant Professors especially Dr. Prashanth , Dr. Tanuja Department of Ophthalmology, Sri Devraj Urs Medical College Tamaka, Kolar for their help and suggestions rendered to me during this study.

I am Immensely thankful to all my PG Colleagues especially Dr. Bhavana, Dr. Tejal, Dr. Iffat, Dr.Poornima , Dr. Shirin for their timely support and encouragement.

My gratitude and thanks to Dr.M.B.Sanikop M.S, (Anatomy), Principal, Sri Devraj Urs Medical College Tamaka, Kolar Sri Devaraj Urs Academy of Higher Education And Research for letting me use the college and hospital facilities and resources.



I would like to thank my fellow postgraduates for their support, cooperation and constant help during the whole study period.

I would like to thank my parents and siblings for their constant encouragement and help.

My heartfelt gratitude to all my patients who submitted themselves most gracefully and wholeheartedly participated in this study.

I sincerely thank my institute Sri Devaraj Urs Medical College, Tamaka, Kolar for giving me a wonderful foundation and forum of knowledge in the field of Ophthalmology which stands for the rest of my life. Last, but not the least, I would like thank to God for reaching the height which I would have never dreamt about without them.

**Date:**

**Signature of the Candidate**

**Time:**

## **LIST OF ABBREVIATIONS USED**

PxS	»	Pseudoexfoliation syndrome
PEX	»	Pseudoexfoliation
IOP	»	Intraocular Pressure
CTR	»	Capsular Tension Ring
CTS	»	Capsular Tension Segment
MSICS	»	Manual Small Incision Cataract surgery
ZD	»	Zonular dehiscence
PCR	»	Posterior capsular rent
VL	»	Vitreous loss
POAG	»	Primary Open Angle Glaucoma
SMC	»	Senile Mature Cataract
SIMC	»	Senile Immature Cataract
SHMC	»	Senile Hyper-mature Cataract

## **ABSTRACT**

### **BACKGROUND**

Pseudoexfoliation syndrome is a common clinically important systemic condition characterized by the pathological production and accumulation of an abnormal fibrillar extracellular material in many intraocular and extra-ocular tissues. Many studies have shown that Pseudoexfoliation syndrome patients have higher rates of intraoperative complications during cataract surgery compared to the patients without the condition.

### **METHODS**

It is a hospital based prospective study of 100 eyes of 100 patients with cataract and Pseudoexfoliation syndrome attending RL Jallapa Hospital and research center ,Tamaka, Kolar

### **OBJECTIVES**

1. Frequency of pseudoexfoliation syndrome
2. To study the complications of cataract surgery in pseudoexfoliation syndrome.

## **RESULTS**

The average age of patients in the study was 63.83 years with a male predominance with equal incidence of unilateral and bilateral involvement with. In the study, 26 (26 %) of the patients had intraoperative complication while 74 (74%) did not. 10 (10 %) of the patients had Zonular dehiscence, 10 (10 %) of the patients had Posterior Capsular Rent and 9 (9 %) of the patients had Vitreous loss. 90 (90 %) of the patients were implanted with intraocular lens after employment of various surgical modifications. 10 (10 %) of the patients were left aphakic due to the above mentioned complications.

## **INTERPRETATION & CONCLUSION**

Inadequate mydriasis is one of the major preoperative complications in eyes with Pseudoexfoliation syndrome which has a bearing on the intraoperative complications. Pupil enlargement procedures are advocated during cataract surgery. Though Small incision cataract surgery in eyes with Pseudoexfoliation syndrome is associated with intraoperative complications ,they can be managed well and good outcome can be expected.

## **KEYWORDS**

Pseudoexfoliation syndrome; Manual Small Incision Cataract Surgery; Intraoperative Complications; Inadequate mydriasis; Intraocular Pressure; Glaucoma; Zonular dialysis; Posterior capsular rent; Vitreous Loss.

## TABLE OF CONTENTS

SL. NO.	NAME	PG. NO.
1.	INTRODUCTION	1
2.	AIMS AND OBJECTIVES	8
3.	REVIEW OF LITERATURE	9
4.	MATERIALS AND METHODS	34
5.	OBSERVATION AND RESULTS	41
6.	DISCUSSION	56
7.	CONCLUSION	67
8.	SUMMARY	69
9.	BIBLIOGRAPHY	72
10.	ANNEXURES	81

## LIST OF TABLES

SL.NO TABLES	PAGE NO.
1. AGE DISTRIBUTION ON PATIENTS WITH PSEUDOEXFOLIATION SYNDROME	40
2. SEX DISTRIBUTION IN PSEUDOEXFOLIATION SYNDROME	41
3. LATERALITY IN EYES WITH PSEUDOEXFOLIATION SYNDROME	42
4. IRIS CHARACTERISTICS IN PSEUDOEXFOLIATION SYNDROME	43
5. ANGLE CONFIGURATION IN PSEUDOEXFOLIATION SYNDROME	44
6. INTRAOCULAR PRESSURE IN PSEUDOEXFOLIATION SYNDROME	45
7. PUPILLARY DILATATION IN PSEUDOEXFOLIATION SYNDROME	46
8. ZONES OF PSEUDOEXFOLIATION ON LENS SURFACE	47
9. TYPE OF CATARACT IN PSEUDOEXFOLIATION SYNDROME	48
10. INTRAOPERATIVE COMPLICATIONS IN PSEUDOEXFOLIATION SYNDROME	49

SL.NO TABLES	PAGE NO.
11. INDIVIDUAL TYPES OF INTRAOPERATIVE COMPLICATIONS IN PATIENTS WITH PSEUDOEXFOLIATION SYNDROME	50
12. SURGICAL OUTCOME IN PATIENTS WITH PSEUDOEXFOLIATION SYNDROME	51
13. INTRAOPERATIVE COMPLICATIONS VERSUS PUPILLARY DILATATION	52
14. SURGICAL MODIFICATIONS IN PATIENTS WITH PSEUDOEXFOLIATION SYNDROME	53

## LIST OF FIGURES

SL.NO FIGURES	PAGE NO.
1. STAGES OF PSEUDOEXFOLIATION SYNDROME	12
2. AGE DISTRIBUTION IN PATIENTS WITH PSEUDOEXFOLIATION SYNDROME	40
3. SEX DISTRIBUTION IN PSEUDOEXFOLIATION SYNDROME	41
4. LATERALITY IN PATIENTS WITH PSEUDOEXFOLIATION SYNDROME	42
5. IRIS CHARACTERISTICS IN PSEUDOEXFOLIATION SYNDROME	43
6. ANGLE CONFIGURATION IN PSEUDOEXFOLIATION SYNDROME	44
7. INTRAOCULAR PRESSURE IN PSEUDOEXFOLIATION SYNDROME	45
8. PUPILLARY DILATATION IN PSEUDOEXFOLIATION SYNDROME	46
9. ZONES OF PSEUDOEXFOLIATION LENS MATERIAL	47
10. TYPE OF CATARACT IN PSEUDOEXFOLIATION SYNDROME	48



SL.NO FIGURES	PAGE NO.
11. INTRAOPERATIVE COMPLICATIONS IN PSEUDOEXFOLIATION SYNDROME	49
12. INDIVIDUAL TYPES OF INTRAOPERATIVE COMPLICATIONS IN PSEUDOEXFOLIATION SYNDROME	50
13. SURGICAL OUTCOME IN PATIENTS WITH PSEUDOEXFOLIATIVE SYNDROME	51
14. INTRAOPERATIVE COMPLICATIONS VERSUS PUPILLARY DILATION	52
15. SURGICAL MODIFICATIONS IN PATIENTS WITH PSEUDOEXFOLIATIVE SYNDROME	53

## LIST OF PHOTOGRAPHS

SL. NO. PHOTOGRAPHS	PAGE NO.
1. PSEUDOEXFOLIATION MATERIAL AT THE PUPILLARY MARGIN AND ANTERIOR SURFACE OF LENS CAPSULE WITH UNDILATED PUPIL.	81
2. ZONE OF PSEUDOEXFOLIATION MATERIAL ON THE ANTERIOR CAPSULE OF LENS AFTER PUPILLARY DILATION.	81
3. MOTH EATEN APPEARANCE IN PSEUDOEXFOLIATION SYNDROME	82
4. HISTOPATHOLOGY OF PSEUDOEXFOLIATION MATERIAL ON THE ANTERIOR LENS CAPSULE.	82
5. SPHINCTEROTOMY CUT AT 10' O CLOCK POSITION. SEEN POSTOPERATIVELY.	83
6. CAPSULAR TENSION RING.	83

# **INTRODUCTION**

---

## INTRODUCTION

Pseudoexfoliation Syndrome is an age related generalized disorder involving abnormal production or turnover of extra-cellular matrix in ocular tissues, orbital tissues, skin and visceral organs.

The exact aetio-pathogenesis of this condition and chemical composition of The material still remains unknown.

Renewed interest in this long known entity results from better awareness of the spectrum of intra-ocular risks not only for open angle glaucoma but also in conjunction with/or intra-ocular surgery, especially cataract extraction.

In the eye, Pseudoexfoliation syndrome is characterized clinically by small white deposits of material in the anterior segment, most commonly in the pupillary border and the anterior Lens capsule. The most consistent diagnostic feature is three distinct zones of pseudoexfoliation material seen on the lens capsule after full dilatation.

1. A translucent, central disc with occasional curled edges.
2. Middle clear zone corresponding to probable contact with the moving iris.
3. Peripheral granular zone, which may have radial striations.

(Central zone is absent in 20% or more cases, but peripheral defect is a consistent finding in all cases. Therefore, pupillary dilatation is a must before lens changes can be seen.)

Additional subtle clinical signs that help in early diagnosis are loss of pigment from peri-pupillary area producing transillumination defects, insufficient mydriasis, and pigment dispersion into anterior chamber after mydriasis, deposition of melanin over trabecular meshwork and Schwalbe's line. The existence of posterior synechiae without any other cause and hemorrhage in the iris stroma after mydriasis are also suggestive of pseudoexfoliation syndrome. Deposition of material on the zonular fibres weakens it leading to phacodonesis, subluxation and dislocation of lens. The presence of secondary open angle glaucoma is known as glaucoma capsulare. The glaucoma has more serious clinical course and worse prognosis than primary open angle glaucoma, often not responding to medical therapy and requiring early surgical intervention. Angle closure glaucoma may also be seen due to pupillary block by forward displaced lens. The corneal endothelium shows decreased cell count and pleomorphism leading to early corneal decompensation at moderate rises in intraocular pressure and after cataract surgery. An increased incidence of nuclear cataract is seen.

Making the diagnosis often requires a careful slit-lamp examination after pupillary dilatation and pseudoexfoliation syndrome frequently goes undiagnosed leading to unexpected problems in management and during surgery.

Due to involvement of virtually all structures by pseudoexfoliation material, patients have a significantly greater risk for a variety of complications during cataract surgery. Poor mydriasis, pigment dispersion, combined with phacodonesis and zonular dialysis predisposes to capsular rupture and vitreous loss. Breakdown of blood-aqueous barrier leads to transient elevations of intraocular pressure and fibrinoid uveitis after surgery. Late complications include

posterior capsular opacification, secondary cataract, and decentration of intra-ocular lens and decompensation of corneal endothelium.

Possible pre-operative and intra-operative measures to avoid or minimize these complications include an increased awareness of pseudoexfoliation syndrome, a careful slit lamp examination after full pupillary dilatation, adequate control of intra-ocular pressure pre-operatively, avoidance of iris manipulation, adequate pupillary dilatation, use of heparin coated intra-ocular lenses and judicious use of steroids post-operatively.

Much remains to be learnt about the pseudoexfoliation material not only at the basic levels of production and by the chemical nature but also with regard to its genetics, epidemiology and treatment. There is an increasing prevalence of pseudoexfoliation syndrome as the mean age of general population increases. Yet the clinical implications of the systemic manifestations of this disorder remain unclear. In view of the multitude of clinical complications, we need to be aware of the risks and specially look for clinical signs of this entity. Pseudoexfoliation syndrome should not be considered as harmless anomaly of the anterior segment but as a potentially catastrophic disease.

## MANIFESTATIONS OF PSEUDOEXFOLIATION SYNDROME

TISSUE INVOLVED		CLINICAL SIGNS
Ocular	lens	Zonular instability, Phacodonesis, Subluxation, Nuclear Cataract.
	Zonules	Zonular instability.
	Iris	Vasculopathy: Blood-aqueous barrier defect,pseudo-uveitis, anterior chamber hypoxia, capillary hemorrhage, iris rigidity, posterior synechiae, poor mydriasis, Asymmetric papillary reaction, stromal / pigment epithelial atrophy, melanin release.
	Trabecular meshwork	Increased resistance to Aqueous outflow, Elevated intra-ocular pressure.
	Cornea	Reduced endothelial cell count. Corneal decompensation Corneal endothelial proliferation.
Extra-ocular	Skin, Extra-ocular muscles, As yet unknown. Heart, Liver, Lung, Kidney, Meninges	

**OCCURRENCE OF COMPLICATIONS OF  
PSEUDOEXFOLIATION SYNDROME**

Spontaneous		Ocular hypertension / glaucoma, Lens Subluxation, Nuclear cataract, Pseudo-uveitis, Corneal endothelial decompensation
Pre-operative ( by medication)	Mydriatics	Poor dilatation, Melanin dispersion, Iris hemorrhage.
	Miotics	Posterior synechiae, Pupillary block, Ciliary block.
Intra-operative	SICS	Poor mydriasis, Zonular dehiscence, PC Rent, Vitreous loss.



Post-operative	General	Blood – aqueous barrier breakdown, Fibrinoid uveitis, Posterior synechiae, Corneal endothelial decompensation.
	SICS	Secondary cataract, Late decentration of Posterior chamber lens, Anterior lens subluxation of lens AC depth irregularity, Conjunctival scarring.

## **AIMS AND OBJECTIVES**

---

## **AIMS AND OBJECTIVES.**

1. Frequency of pseudoexfoliation syndrome
2. To study the complications of cataract surgery in pseudoexfoliation syndrome.

## **REVIEW OF LITERATURE**

---

## REVIEW OF LITERATURE

In 1917, **Lindberg**<sup>1</sup> described grayish or bluish flakes of material on the pupillary border in some patients with glaucoma. **Vogt**<sup>2</sup> later hypothesized that this material represented degenerative changes of the lens capsule followed by secondary desquamation and proposed the term senile exfoliation of the lens capsule. **Busacca**<sup>3</sup> argued that the exfoliative material represented deposition of material formed elsewhere in the eye rather than degenerative changes of the lens capsule. **Dvorak-Theobald**<sup>4</sup> subsequently showed that exfoliative material differed histochemically from lens capsule and, to differentiate this condition from true exfoliation of the lens capsule secondary to infrared exposure, suggested the term pseudoexfoliation of the lens capsule. Subsequent electron microscopic studies by **Ashton**<sup>5</sup> and associates and **Bertelsen**<sup>6</sup> and coworkers indicate that the anterior lens capsule was directly affected in this disorder. Bertelsen and associates suggest that pre-equatorial lens epithelial cells produced the abnormal fibrillar substance and recommend the term fibrillopathia epitheliocapsularis. **Eagle and colleagues**<sup>7</sup>, who believe that the material represented abnormal basement membrane secretions, have called this condition basement membrane exfoliation syndrome. The terms exfoliation syndrome and pseudoexfoliation syndrome are now most commonly used to designate this disorder and are used interchangeably in current literature. However, since recent ultrastructural studies indicate that the material on the lens capsule is derived, at least in part, from the lens, it is proposed that the disorder be called exfoliation syndrome (XFS).<sup>8-10</sup>

## **EPIDEMIOLOGY**

Pseudoexfoliation Syndrome is of global distribution. The reported prevalence of Pseudoexfoliation Syndrome both with and without glaucoma has varied widely. This reflects a combination of true difference due to racial, ethnic or as yet unknown factors; the clinical criteria used to detect early stages and/or more subtle changes, the method and thoroughness of examination and awareness of the examiner. In US population, the **Framingham Eye Study**<sup>11</sup> revealed the overall prevalence of Pseudoexfoliation syndrome to be 0.6% in 52 – 64 years old, rising to 5% in 75 – 85 years old. In India, the prevalence rates reported were 1.88% **Sood N.N.**<sup>12</sup> (1965), 7.4% **Lamba and Giridhar**<sup>13</sup> (1984). The prevalence rate in south India is 3.8% **Aravind H et al**<sup>14</sup> (2003) . In a given population, the actual prevalence of Pseudoexfoliation Syndrome is probably twice that which is visible on clinical examination. Many cases go undetected because of failure to dilate the pupil or to examine the lens with the slit lamp after dilatation of the pupil.

The prevalence increases with age, the disease most commonly manifesting between 60–70 years. But Pseudoexfoliation Syndrome might well be a condition that starts in mid-adulthood but becomes frankly manifest only in later years. Sex ratio reports are conflicting.

A hereditary transmission of Pseudoexfoliation Syndrome is not yet clarified. **Tarkkanen**<sup>15</sup> (1962) suggested the presence of a gene bearing 3 characteristics, an abnormality of the drainage channels of the aqueous, Pseudoexfoliation and degeneration of the pigment epithelium of the iris. Variations in the expressivity of this gene would explain why the 3 events are sometimes found together and why sometimes only 1 or 2 is present.

**Kelvin Y.C. Lee et al**<sup>16</sup> studied about XFS/XFG associations with polymorphisms with R141L, G153D and intronic located in the 1<sup>st</sup> exon of the lysyl oxidase like 1 gene (LOXL1) on Chromosome 15q 21 (1). Asian populations including Indians reported associations with LOXL1 and XFS.

**R.R. Allingham et al**<sup>17</sup> (2001) investigated 6 islandic families each of which had at least 1 member affected by Pseudoexfoliation Syndrome they concluded that Ps Pseudoexfoliation Syndrome is an inherited condition with transmission to the 2<sup>nd</sup> generation through an affected mother.

There are no unequivocal findings regarding the role of environmental factors in the development of Pseudoexfoliation Syndrome.

It is now known that Pseudoexfoliation Syndrome is essentially a bilateral condition and unilateral cases only represent an earlier period in the natural history of the condition. When only 1 eye is involved clinically, the other eye often has abnormal aqueous humour dynamics or glaucomatous damage.<sup>11</sup>

## **CLINICAL FEATURES**

### **1. OCULAR MANIFESTATIONS**<sup>18, 19, 20, 21, 22, 23, 24</sup>

#### **a) LENS AND ZONULES**

Deposits of white flaky material on the anterior lens surface are the most consistent and important diagnostic of Pseudoexfoliation Syndrome. The classic pattern consists of 3 distinct zones that become visible when the pupil is fully dilated – a relatively homogeneous central disk corresponding roughly to the diameter of the pupil, a granular often layered peripheral zone and a clear area separating the 2. The central zone is homogeneous white sheet lying on the anterior pole of the lens capsule. Its diameter varies between 1.5 – 3 mm and it is usually slightly smaller than

the physiological pupil. The edges of the disk are often rolled equatorially. The central disk is absent in 20 – 60% of cases. It is often initially overlooked but with careful examination after dilatation, a subtle area of Pseudoexfoliation material may be noted especially when compared to the adjacent intermediate clear zone. It may be granular in the periphery and frosty white centrally and radial striations are often seen. It may be layered. Axially it is bounded partly by curled edges and partly by tongue shaped projections. Equatorially it extends as granular tongue shaped projections which merge into the normal capsule before reaching the anterior zone of insertion of the zonular fibres. The peripheral band may be situated close to the equator in some eyes and more axially in others. The granularity of the peripheral layers is consistent with undisturbed accumulation of Pseudoexfoliation material. Whereas the classical picture of Pseudoexfoliation Syndrome has often been described, the early stages have not been well defined. A precursor of Pseudoexfoliation material is thought to be initially deposited diffusely on the lens surface. A homogeneous “ground glass” or “matte” appearance of the lens surface in one eye compared to the other may represent a very early (pre-capsular) stage. In a perhaps slightly later (pre-granular) stage, there may be very faint radiant non-granular striae on middle third of the anterior capsule behind the iris. Ultra structurally, the pre-capsular layer at this stage consists of micro-fibrils, but not mature exfoliation fibrils. To visualize the earlier stages at the slit lamp, placing the slit beam at 45° to the axis of observation reducing the light source and focusing temporarily 2 – 3 mm from the centre of the lens may help to highlight the subtle deposits on the lens surface. The intermediate clear zone is created by rubbing of the iris over the surface of the lens during pupillary movement. As the pre-capsular layer becomes thicker the iris sphincter begins to rub against it during normal pupillary




movement. Faint clefts begin to form where Pseudoexfoliation material is rubbed away in what will eventually become the clear zone. With time, these clefts increase in size and begin to become confluent. Eventually only small bridges may remain as an indication of the previous layer of Pseudoexfoliation material in the intermediate zone. In some patients the central disk may become thick enough to peel away in sheets from the lens, as may the peripheral zone, giving rise to appearance of True Exfoliation Syndrome. Chronic pupillary dilatation also permits undisturbed accumulation of Pseudoexfoliation material. Clinical classification of various stages is based mainly but not only on the findings of the anterior lens capsule.<sup>18</sup>

→ SUSPECT PSEUDOEXFOLIATION SYNDROME:

- Early Pseudoexfoliation Syndrome (Electron Microscopy): Pre-capsular layer.
- Masked/Suspected Pseudoexfoliation Syndrome:
  - Posterior synechiae without any obvious cause.

→ DEFINITE PSEUDOEXFOLIATION SYNDROME:

- Mini-Pseudoexfoliation Syndrome: Focal defects in pre-capsular layer especially supero-nasally.
- Classic Pseudoexfoliation Syndrome: Late stage.



Phacodonesis is common but not always associated with iridodonesis, perhaps attributable to increased iris rigidity.

Spontaneous subluxation and dislocation of lens can occur; the denser the Pseudoexfoliation material, the more likely there is to be phacodonesis. Lens dislocation is more common inferiorly.

The Zonular fibrils coated with varying amounts of Pseudoexfoliation material become stretched and eventually break. Break is not seen to occur from the attachment to the zonular lamellae but at their ciliary attachments. The broken fibers may be seen waving gently in the aqueous. Subsequently the fibers become shorter and thicker and finally appearing as irregular clumps on the lens surface. The fibers that break 1<sup>st</sup> are those behind the equator and those just anterior to the equator

remain intact the longest.

#### b) IRIS AND PUPIL

Next to lens, Pseudoexfoliation material is most prominent at the pupillary border. It may be extensive or minimal. The iris is more rigid because of the material. Pigment loss from the iris sphincter region and its deposition on the anterior chamber structures is the hallmark of Pseudoexfoliation Syndrome. The material on the lens causes rupture of iris pigment epithelial cells at the ruff and sphincter region with concomitant dispersion of pigment into anterior chamber. Loss of iris pigment and its deposition throughout the anterior segment are reflected in iris sphincter region transillumination, loss of pupillary ruff, increased trabecular pigmentation and pigment deposition on the iris surface. Extensive depigmentation may be noted over the entire sphincter region, which appears as a diffuse starry sky pattern on transillumination or moth eaten appearance.

Pseudoexfoliation Syndrome predisposes to formation of synechiae between iris pigment epithelium and the anterior lens capsule. Posterior synechiae are more prone to form between the iris and intra-ocular lens post operatively. Iris blood vessel

abnormalities include narrow or obliterate lumen, with marked alteration of iris vasculature, vessel dropout with collateral formation and iris hypo perfusion leading to patchy iris neo-vascularization.<sup>19</sup>

Inflammation after cataract extraction is more common and a transient fibrinoid reaction attributed to breakdown of Blood-aqueous barrier may occur.<sup>20</sup>

Intra-stromal hemorrhage after mydriasis is indicative of vascular damage. Atrophic changes of sphincter and dilator muscle tissues, possibly because of hypoxia, and apparent impairment of muscle cells by Pseudoexfoliation material may contribute to poor pupillary dilatation. Reduction of stromal elasticity by accumulating Pseudoexfoliation material may also play a role in poor mydriasis. Dispersion of melanin granules after diagnostic mydriasis or surgery can be so pronounced that heterochromia iridium may be produced. The mechanism of melanin liberation is related to degenerative changes and cell membrane ruptures of the posterior pigmented epithelial cells due to extra-cellular Pseudoexfoliation material. Marked intra-ocular pressure rise after mydriasis correlates with the amount of the pigment liberated.

#### c) CILLIARY BODY

The ciliary processes were examined clinically by **Mizuno and Muroi**<sup>21</sup> with special type of Gonioscopy lens, almost all eyes with exfoliation showed accumulation of material on the zonules and ciliary body.

#### d) GLAUCOMA AND PSEUDOEXFOLIATION SYNDROME<sup>22,23</sup>:

While the existence of association between Pseudoexfoliation Syndrome and Open Angle Glaucoma has been well known, the mechanisms are still not clarified. There is an increase in the aqueous outflow resistance probably due to trabecular cell

dysfunction, blockage of meshwork by Pseudoexfoliation Syndrome liberated pigment and concomitant primary open angle glaucoma.

In patients with pseudoexfoliation syndrome, 20% have glaucoma and increased IOP at the time of diagnosis. Patients who have pseudoexfoliation syndrome but not glaucoma should be considered vulnerable to glaucoma, because 15% of such patients develop increased IOP within 10 years. This underscores the need for careful follow-up in patients who have pseudoexfoliation syndrome. Pseudoexfoliation syndrome accounts for 15-20% of cases of open angle glaucoma.

Glaucoma in Pseudoexfoliation Syndrome has a more serious clinical course and worse prognosis than Primary Open Angle Glaucoma. There is a significantly higher frequency and severity of optic nerve damage at the time of diagnosis, worse visual field damage, and poorer response to medications, more severe clinical course and more frequent necessity of surgical interventions. In normotensive eyes, with Pseudoexfoliation Syndrome the mean Intra-ocular pressure is higher than in eyes without Pseudoexfoliation Syndrome. In patients with elevated Intra-ocular pressure, mean intra-ocular pressure is higher at the time of diagnosis in patients with Pseudoexfoliation Syndrome, than in those with primary open angle glaucoma. Glaucomatous damage at the time of diagnosis is more severe and progression is also more rapid in eyes with Pseudoexfoliative glaucoma.

A number of characteristics predispose to development of angle closure glaucoma in eyes with Pseudoexfoliation Syndrome. Pupillary block may be caused by combination of posterior synechiae, increased iris thickness or rigidity or anterior lens movement secondary to zonular weakness or dialysis.

#### e) ANGLE CHARACTERISTICS<sup>22</sup>:

As the iris is more rigid than normal, aqueous pressure in the posterior chamber causes it to bulge at the weakest point which is the iris root. Thereby, the localized iris bombe near the iris root narrows the angle, giving a pseudo-plateau iris configuration on gonioscopy and leads to chronic angle closure glaucoma.

Increased trabecular pigmentation is a prominent sign and is apparent in virtually all patients with clinically evident disease. The pigment is splotchy and less well defined. It is an early diagnostic finding preceding appearance of Pseudoexfoliation material on the pupillary margin and the anterior lens capsule. It is almost always dense in the involved eye and increases in eyes with Pseudoexfoliative glaucoma. The degree of pigmentation correlates with elevated intra-ocular pressure. Pigment on Schwalbe's line is seen as a wavy line known as Sampolesi's Line which is also an early sign of Pseudoexfoliation Syndrome.

#### f) VITREOUS

Vitreous changes commonly accompany Pseudoexfoliation Syndrome since hyaluronic acid and Pseudoexfoliation material are both acid mucopolysaccharides. A change in composition of aqueous in Pseudoexfoliation Syndrome could derange metabolism of hyalocytes leading to impaired production of hyaluronic acid and liquefaction.

#### g) CONJUNCTIVA AND CORNEA:

Clinically the Conjunctiva is normal. However, fluorescein angiography reveals loss of regular limbal vascular pattern and areas of neovascularisation and congestion of anterior ciliary vessels. Scattered flakes of Pseudoexfoliation material may be observed on the endothelial surface of the cornea. Specular microscopy demonstrates a significantly reduced endothelial cell density even with normal intra-ocular

pressure, together with morphological changes in size and shape of the endothelial cells in both affected eyes and un-involved fellow eyes. Decreased endothelial cell density does not necessarily correlate with the severity of glaucoma but it has been correlated with the extent of pigment dispersion. Central corneal thickness is increased reflecting early corneal dysfunction. These changes may help in early diagnosis and in pre-operative assessment prior to cataract extraction. These eyes can develop early corneal endothelial decompensation at only moderate rises of intra-ocular pressure or after cataract surgery.

## 2. SYSTEMIC MANIFESTATIONS<sup>23, 25, 26</sup>

Ultrastructural studies performed on eyes during autopsy suggest that Pseudoexfoliation syndrome is a multisystem disorder, Pseudoexfoliation material has been found in a number of organs, which include skin, lungs, gallbladder, liver, myocardium, kidney, bladder and Meninges. Associations of aneurysms of abdominal aorta and Pseudoexfoliation syndrome have been extensively studied. The staining of the material in these organs is positive for elastin and human amyloid P protein, which is similar to the staining pattern characteristic of the material found in the eye. These findings provide evidence for the systemic nature of Pseudoexfoliation syndrome, which involves an aberrant connective tissue metabolism throughout the body. Patients with pseudoexfoliation syndrome are found to have sensory neural-deafness.

## **THEORIES ON ORIGIN OF PSEUDOEXFOLIATION MATERIAL**

### 1. BASEMENT MEMBRANE THEORY:

There is extensive support for the hypothesis that pseudoexfoliation syndrome represents a disorder of extra-cellular matrix characterized by overproduction or abnormal breakdown of cell surface associated material, the biochemical nature of

which remains unclear. With the advent of the electron microscope, extensive studies on the pseudoexfoliation material were done and its origin was ascribed to be basement membrane of the lens capsule, iris, ciliary body and conjunctiva. **Schlotzer-Schrehardt et al**<sup>25</sup> in 1992 confirmed systemic involvement of the viscera by pseudoexfoliation material using a transmission electron microscopy. Typical pseudoexfoliation fibers were identified in autopsy tissue specimens of skin, heart, lungs, liver, kidney and cerebral meninges in addition to the classic intraocular locations leading to the term pseudoexfoliation syndrome.

The production of the exfoliation material may be related to disordered basement membrane metabolism and **Harnisch et al**<sup>27</sup> in 1981 using the indirect immunoperoxidase method, found that the fibrils contained a basement membrane proteoglycan. Anti-basement membrane proteoglycan antibodies to lens material reacted strongly with exfoliation material, implicating lens epithelium and its production.

## 2. ELASTIC MICRO-FIBRAL THEORY

Since exfoliation material is immunologically related to elastic tissue, Li et al in 1989 proposed that exfoliation fibers have peripheral binding sites for Amyloid P protein similar to those present on normal elastic fibers. There are histochemical and antigenic similarities between zonular elastic microfibrils and exfoliation material. **Garner and Alexander**<sup>28</sup> in 1984 suggested that Oxytalan, a micro-fibrillar component of elastic tissue present in the body in areas of mechanical stress is a constituent of the exfoliation fibrils. **Roh et al** in 1987 found mature and intermediate micro-fibrils adjacent to fibroblasts in close proximity to elastic tissue in the conjunctiva. **Streeten et al**<sup>29</sup> in 1987 found histochemical similarities between zonular elastic micro-fibrils and pseudoexfoliation material and a resemblance of the



larger micro-fibrils of a ground substance to zonular and other oxytalan micro-fibrils. The strong anatomic association between pseudoexfoliation fibers with elastosis in conjunctival specimens led the authors to suggest that pseudoexfoliation fibers themselves might be a form of elastosis, possibly resulting from abnormal aggregation of components related to elastic micro-fibrils. Elastin and elastic micro-fibril protein were demonstrated in pseudoexfoliation material – their production might reflect an abnormal stimulus or defective regulation of matrix synthesis. **Schlotzer – Schrehardt et al**<sup>30</sup> in 1998 analyzed by electron microscopy the matrix of the pseudoexfoliation material and demonstrated in to be fibrilin positive fibers, supporting the elastic micro fibril theory of its production.

### 3. AMYLOID THEORY

**Repo L.P. Naucharinen et al**<sup>31</sup> in 1996 examined by light and electron microscopy 13 biopsy specimens of iris tissue from patients with pseudoexfoliation syndrome undergoing cataract surgery. They showed that pseudoexfoliation material is associated with amyloid and in some eyes; miosis is associated with degenerative changes, both in stromal tissue and in muscular layers of the iris.

**Tsukahara and Matsuo**<sup>32</sup> described patients with both primary familial amyloidosis and exfoliation.

### 4. LYSOZOMAL THEORY

**Mizuno et al**<sup>33</sup> in 1980 found histochemical evidence of high acid phosphatase activity, suggesting that lysozymes were involved in the production of exfoliation material. Possible rupture of pigment epithelial cells may account for lyzosomal involvement. Proteolytic enzymes present in lysosomes may facilitate granular disintegration.

**Baba**<sup>34</sup> in 1982 demonstrated a lipoprotein in exfoliation material and felt that this might be the result of the high permeability of vessels in the anterior segment. He also found that material was a sulphated glycosaminoglycan and suggested that abnormal glycosaminoglycan metabolism precedes the formation of the material.

Immunochemical studies have revealed heparin sulphate, chondroitin sulphate proteoglycans, laminin, entactin/nidogen, fibronectin and amyloid P protein to be integral constituents of exfoliation material. Type IV collagen is restricted to a microfibrillar layer interposed between the capsular surface and typical exfoliation material. Type IV collagen mediates cell attachment and might be instrumental in adherence of exfoliation material to the anterior central capsule. The additional presence of elastin epitopes indicate that exfoliation material is a multi-component expression of a disordered extracellular matrix synthesis, including the incorporation of the principle noncollagenous basement membrane components. Extensive labeling of exfoliation material for chondroitin sulphate suggests an over-production and abnormal production of glycosaminoglycans to be one of the key changes in this disorder. Exfoliation material contains but does not represent true basement membrane material because of absence of Type IV collagen and the additional presence of elastin epitopes.

Transmission electron microscopy and high resolution scanning electron microscopy and demonstrated pseudoexfoliation material to contain keratan and dermatan sulphate. They postulated that pseudoexfoliative material was produced due to abnormality in proteoglycans.

None of the histochemical or enzymatic studies have succeeded in elucidating the exact source of pseudoexfoliation material. This along with the increased chances

of surgical complications continues to arouse great interest in pseudoexfoliation syndrome.

### **STRUCTURE OF PSEUDOEXFOLIATION MATERIAL**

The Pseudoexfoliation Material consists of an irregular meshwork of randomly oriented cross-banded fibrils measuring about 30 nm in diameter within a loose fibro-granular matrix containing 6 – 10 nm micro fibrils.

**Davanger**<sup>35, 36</sup>, (1978) described the fibrils as consisting of a protein core surrounded by polysaccharide side chains. The fibrils are formed from lateral aggregations of filaments.

The abnormally produced Pseudoexfoliation Material on light microscopy is a PAS positive, eosinophilic brush like nodular or feathery aggregate. On scanning electron microscopy these aggregates are composed of an irregular tangle of fibrils.

The fibrils are intermingled with normal micro-fibrils and are embedded in an amorphous inter-fibrillar ground substance, most probably glycosaminoglycans. The extra-ocular Pseudoexfoliation Material is similar except that there is more matrix and less distinct banding pattern.

Indirect histochemical and immune histochemical evidence suggests a complex glycoprotein/proteoglycan like structure composed of a protein core surrounded by glycol-conjugates probably glycosaminoglycans forming the amorphous substance.

### **CATARACT SURGERY IN PSEUDOEXFOLIATION SYNDROME**

Patients with Pseudoexfoliation Syndrome are much more prone to have complications at the time of cataract extraction. Eyes with Pseudoexfoliation

Syndrome dilate less well and have greater incidence of capsular rupture, zonular dehiscence and vitreous loss. Pupillary diameter and zonular fragility have been suggested as the most important risk factors for capsular rupture and vitreous loss. The presence of phacodonesis has been related to poor mydriasis, cataract, and presence of glaucoma and trabecular pigmentation, all a reflection of the severity of involvement and should serve a warning sign. A shallow anterior chamber may indicate zonular instability. Post-operatively, transient intra-ocular pressure elevations are common. Posterior capsular opacification is more common. Late postoperative decentration of intra-ocular lens and capsular bag are common, and is related to zonular weakness. Capsular contraction syndrome if exaggerated, can lead to intra-ocular lens dislocation. Secondary cataract is more common because of aggravated blood-aqueous barrier breakdown.

**Skuta G. L., Parrish R. K. et al**<sup>37</sup> (1987) showed an increased incidence of zonular dialysis in patients with Pseudoexfoliation Syndrome during cataract surgery. They stated that pre-operative phacodonesis, anterior chamber depth asymmetry and excessive lens movement during anterior capsulotomy should alert to the presence of zonular dialysis.

**Naumann G. O., Kuchle M. Schonherr U**<sup>38</sup> (1989) noted a seven fold increase in vitreous loss in 72 patients with Pseudoexfoliation Syndrome undergoing cataract surgery.

**Wang L., Yamasita R. et al**<sup>39</sup> (1999) studied 26 eyes with Pseudoexfoliation Syndrome with specular microscopy and quantified the aqueous flare with laser flare cell meter. They showed that the corneal endothelial cell density was significantly decreased in eyes with Pseudoexfoliation Syndrome and an inverse correlation was

shown with the flare. The authors concluded that a decrease in the endothelial cells may correlate with a disruption of blood-aqueous barrier.

**Kuchle M, Naumann.H et al<sup>40</sup>** (1997) emphasized the fact that pseudoexfoliation syndrome is frequently associated with impairment of blood-aqueous barrier and thereby have higher frequency of secondary cataract post cataract surgery.

**Lumme P. Lattikaanen L<sup>41</sup>** (1993) performed a prospective study of 351 patients undergoing cataract surgery. In their study the prevalence of Pseudoexfoliation Syndrome was more in patients greater than 70 years. Pseudoexfoliation Syndrome increased the risk on intra-operative complications either directly (rupture of zonules) or through poor dilation of pupil (rupture of posterior lens capsule). The occurrence of vitreous loss was four fold and the need to use anterior chamber intra-ocular lens was tenfold in these patients.

**Moreno et al<sup>42</sup>** (2000) suggested irido-phacodonesis, poor dilatation and presence of glaucoma as the clinical factors related to capsular rupture during cataract surgery.

**Freyler H. Radax U<sup>43</sup>** (1994) compared Extra-capsular cataract surgery with phacoemulsification in 311 and 68 patients respectively. Miosis and phacodonesis were reported as the primary risk factors for cataract surgery associated with Pseudoexfoliation Syndrome. Compared with extra-capsular cataract surgery, phacoemulsification had significantly fewer complications with regard to miosis but not phacodonesis. They advised applying a small iris retractor, hooks to stem the complications arising from miosis.

**Stanila A<sup>44</sup>** (1996) noted that out of 868 patients undergoing cataract surgery, 10% had Pseudoexfoliation Syndrome and these patients had an increased incidence

of insufficient dilatation of pupil, posterior capsular tears, Vitreous loss, increase in post-operative intra-ocular pressure and more frequent opacification of posterior capsule.

**Kuchle et al**<sup>45</sup> (2000) suggested that a shallow anterior chamber depth of less than 2.5 mm pre-operatively was indicative of zonular instability and should alert the surgeon of intra-operative complications.

**Bayramlar et al**<sup>46</sup> (2007), conducted a retrospective study in 225 eyes of 187 patients, of which 99 eyes had pseudoexfoliation syndrome. Preoperative data collected were cataract maturity level, best corrected visual acuity and intraoperative posterior capsule complications. In this study, he interpreted that in manual small incision cataract surgery, pseudoexfoliation syndrome has an increased intraoperative posterior capsule complication rate that increases at the level of cataract maturity increases and the preoperative visual acuity decreases.

**Albert Galand MD, Michael Kuchle MD, Etienne Thchet MD**<sup>47</sup> (2004) at a symposium held during the 21st congress of the ESCRS reviewed the pathophysiological alterations associated with pseudoexfoliation, the consequences of cataract surgery, and the considerations for surgical modifications and intraocular lens selection. They stressed on poor mydriasis, a prominent feature of pseudoexfoliative eyes and its management by injection of high viscosity viscoelastic agent, also advocated use of iris hooks, either plastic or metallic as necessary. Dr. Hachet cautioned against performing sphincterotomy, which resulted in persistent dilatation and poor postoperative chemosis, he recommended against use of circular plastic dilator to push the pupil rim aside, also advocated on use of capsular tension rings. Foldable intraocular lens is desirable to minimize the induction of blood-aqueous barrier breakdown and the accompanying increased risks for postoperative

complications. Also hydrophobic acrylic and silicone are associated with a low rate of posterior capsular opacification, but hydrophobic acrylic has an additional advantage as it causes the least amount of capsular contraction. For haptics, overall, open loop haptics are probably preferred, and PMMA may be better than prolene. Dr Kuchle also discouraged the use of plate haptic design or accommodative intraocular lens in patients with pseudoexfoliation.

**Vickie Lee and Anthony Maloof**<sup>48</sup> (2002) stated that a CTR allows for the expansion and stabilization of the capsular bag by redistributing forces with the resulting tautness of bag providing counter-traction to facilitate cataract surgery and cortical aspiration. This is extremely useful for moderate degrees (i.e. up to 5 hours) of zonular dialysis. The CTR can be inserted after the completion of capsulorrhexis but before hydrodissection.

**Howard Fine**<sup>49</sup> (2008) CTR will convert a high risk case into a routine case when there is compromised zonular integrity. CTR work because the ring diameter is larger than the capsule diameter so that there is centrifugal force on the capsular fornix and this distributes focal forces. Any focal force on the capsule cannot be transmitted only to the adjacent zonules with an unzipping of the zonular apparatus – the ring makes that focal force distributed circumferentially to the entire zonular apparatus. In cases of advanced zonulopathy with overt subluxation of the capsular bag, the capsular tension segments (CTS) can be used instead of CTR. The CTS is a 120° partial CTR that features an islet positioned within the capsulorrhexis that can receive an iris hook for support. Two CTS can be used to support a very loose bag. The CTS can also be used in eyes with anterior or posterior capsular tears. The CTS are also designed for suture sclera fixation, for long term capsular bag centration.

## **MANAGEMENT OF CATARACT SURGERY IN PSEUDOEXFOLIATION SYNDROME<sup>50, 51, 52, 53, 54</sup>**

These are several important points to remember for cataract surgery in eyes with Pseudoexfoliation syndrome.

### **1. MAKING THE DIAGNOSIS**

Limited pharmacological mydriasis can adversely affect the ability to make the diagnosis. Flaky deposits on the corneal endothelium is one, clue in assessing the condition. This material can be differentiated from true keratic precipitate by their bright white color and fluffy appearance. When differentiation is difficult, a one to two week course of topical steroids can aid in diagnosis, as keratic precipitates change in appearance or location or disappear with topical steroid use but have no effect on Pseudoexfoliation material.

An unusually shallow anterior chamber depth from zonular instability can indicate Pseudoexfoliation especially if it is asymmetrical. Even though a patient's cataract and symptomatic complaints are monocular, the contralateral eye may have subtle findings of Pseudoexfoliation which may not be seen in the planned surgical eye. Even if Pseudoexfoliation material is not clinically visible on the corneal endothelium, the cell count may be significantly reduced and the cells that remain may not function well, hence additional endothelial protection including a "pseudoplastic" viscoelastic such as healon is advised.

### **2. MAXIMAL DILATATION OF PUPIL DURING SURGERY.**

Poor mydriasis, a well known feature of Pseudoexfoliation syndrome can seriously hamper the surgeon's view, additional pupillary dilatation may also be necessary. Several mechanical means can temporarily dilate the pupil during surgery. These include flexible iris retractors, titanium iris retractors, flexible pupil dilating



rings and rigid dilating rings. Pupil stretching maneuvers like sphincterotomies are an inexpensive and easier alternative. While these are effective, excessive inflammatory responses due to the compromised blood-aqueous barrier in these eyes are well documented. Further, the iris is more flaccid in Pseudoexfoliation syndrome and more likely to be inadvertently aspirated; mechanical means to augment mydriasis is to also keep the floppy iris margin away from the aspiration port or cannula. Care should be taken to avoid excessive iris trauma and over-inflation of the anterior chamber with viscoelastic, which can cause posterior pressure on the lens and can further damage the weakened zonules.

### 3. ENSURING ADEQUATE CAPSULORRHEXIS/CAPSULOTOMY.

Capsulorrhexis/capsulotomy creation is more difficult in these cases, as there is no counter-traction during tearing of the anterior lens capsule. This can present as a star pattern of capsular folds radiating from your instrument when piercing the anterior lens capsule and as wrinkling and looseness of the capsule.

The solution, as described by Thomas Neuhann, MD, of Germany, is to provide counter-traction via the non-dominant hand using a chopper or other second instrument via the paracentesis, while using the dominant hand to perform the capsulorrhexis via the main incision. Because of the tendency for anterior capsular phimosis and further zonular stress, a large capsulorrhexis should be performed, at least 5.5 mm in diameter. Staining the capsule with indocyanine green or trypan blue is useful. The Pseudoexfoliation material has a higher affinity for indocyanine green stain than unaffected capsule.

### 4. ATTENTION TO PHACODONESIS WHILE PERFORMING CAPSULORRHEXIS/CAPSULOTOMY.

Weak zonules is one of the most notorious, common and significant problem faced by cataract surgeon in Pseudoexfoliation syndrome. The degree of weakening though highly variable appears to increase with apparently increasing amount of deposits. Dislocation of the nucleus into the vitreous cavity may occur even during routine hydrodissection. During capsulorrhexis or capsulotomy creation, diffuse zonular weakness or laxity may be sensed. Once this weakness is apparent, the risk of creating zonular dialysis looms large. In such cases, flexible “iris” retractors can engage the capsulorrhexis margin and stabilize the loosened capsular bag.

#### 5. MANAGEMENT OF ZONULAR DIALYSIS.

If a small or moderate zonular dehiscence occurs, a standard capsular tension ring can re-expand the capsular bag and redistribute the mechanical stresses evenly across the remaining zonules. The capsular tension ring (CTR) can be manually implanted into the fornix of the capsular bag or injected with the inserter device. For a large zonular dehiscence, a suture-fixated, modified Cionni ring with one or two fixation eyelets will re-expand the capsular bag and secure the capsular bag or intraocular lens complex to the sclera wall.

#### 6. CHOICE OF INTRAOCULAR LENS.

Capsular contraction is more likely since there is reduced zonular counter-traction against the centripetal forces of the remaining lens epithelial cells. Capsulorrhexis of 5mm or greater, and use of a capsular tension ring to reduce the risk of this complication is advisable. As capsular contraction is more common with silicone intraocular lens, another material is preferred. An intraocular lens with a sharp posterior edge to reduce lens epithelial cell migration and subsequent posterior capsular opacification is recommended. Pseudoexfoliation syndrome adds to the challenges of cataract surgery.

Some of these challenges are significant. With the use of dyes, capsule retractors and implant rings and meticulous attention to surgical technique, cataract surgery in Pseudoexfoliation syndrome may be safely performed.

## **MATERIALS AND METHODS**

---

## **MATERIALS AND METHODS.**

### **TITLE OF THE STUDY:**

“Prospective Two Years Study Of Complications Of Manual Small Incision Cataract Surgery In Eyes With Pseudoexfoliation Syndrome”.

**SOURCE OF DATA:** Patients admitted with cataract and pseudoexfoliation at R.L.J. HOSPITAL AND RESEARCH CENTRE,TAMAKA, KOLAR attached to SRI DEVRAJ URS MEDICAL COLLEGE between November 2008 and July 2010 were prospectively analysed. Total number of approximately 100 cases of acquired cataract fulfilling the selection criteria were included in the study after their informed consent.

### **SAMPLE SIZE:**

A total number of 100 eyes of 100 patients were selected for the study that had cataract in patients with pseudoexfoliation syndrome.

### **INCLUSION CRITERIA:**

- Patients undergoing manual small incision cataract surgery with pseudoexfoliation syndrome
- Patients with pseudoexfoliation syndrome as evidenced by pseudoexfoliation material on the pupillary margin and on the anterior surface of lens capsule.
- Patients with senile cataracts –immature, mature, hypermature cataracts in pseudoexfoliation syndrome.

### **EXCLUSION CRITERIA:**

- Traumatic cataracts.
- Complicated cataracts.
- Subluxated lens without pseudoexfoliation syndrome.

- Patients with senile cataracts – immature, mature, hypermature cataracts without pseudoexfoliation syndrome.
- Patients posted for other surgeries other than manual small incision cataract surgery in patients with pseudoexfoliation syndrome .

**PREOPERATIVE EVALUATION.**

1. Visual acuity testing for distance and near using Snellen’s distant chart and near vision chart respectively.
2. Refraction and correction where required.
3. External ocular examination.
4. Slit lamp biomicroscopic examination for evidence of the following findings.
  - Pseudoexfoliation material in the pupillary margins.
  - Moth eaten appearance of the iris.
  - Morphological alterations of the cornea
  - Anterior chamber depth and pigment dispersion in the anterior chamber
  - Iridodonesis.
  - Presence of posterior synechiae.
  - Zones of Pseudoexfoliation on the anterior surface of the lens capsule.
  - Phacodonesis or frank subluxation/dislocation of lens.
  - Measurement of pupil size before and after dilatation of pupil.
  - Pupillary reactions.
5. Tonometry using Schiottz tonometer.
6. Gonioscopy with Goldmann three mirror lens in all patients with pseudoexfoliation syndrome. The following points were specifically evaluated.

- The extent of trabecular pigmentation which was graded as:

- Grade 0 → Nil
- Grade 1 → Faint Pigmentation
- Grade 2 → Average Pigmentation
- Grade 3 → Moderate Pigmentation
- Grade 4 → Heavy Pigmentation

- The presence of pseudoexfoliation material in the angle.
- The presence of Sampolesi's line.
- The grading of angle width according to Shaffer's grading.

<b>Grade</b>	<b>Angle width (degree)</b>	<b>configuration</b>	<b>Chance of closure</b>	<b>Structure visible on gonioscopy.</b>
4	35-45	Wide open	Nil	From Schwalbe's line to ciliary body
3	20-35	Open	Nil	From Schwalbe's line to sclera spur
2	20	Moderately Narrow	Possible	From Schwalbe's line to Trabecular meshwork
1	10	Very narrow	High	Schwalbe's line only
0	0	Closed	Closed	None of the structures visible.

- The pupils were then dilated with a combination of 5% phenylephrine and tropicamide 0.8% drop was instilled every 5 minutes for 15 minutes interval.

8. This was followed by slit lamp examination for
  - Measuring pupil size.
  - Examination of lens capsule for central and peripheral zones of pseudoexfoliation material deposition.
  - Evaluation of lens for the type of cataract.
9. Fundoscopy
10. Lacrimal patency test
11. Keratometry
12. A-scan and Intraocular lens power calculation by SRK-2 formula.

Other investigations included

- Urine examination for detection of sugar and albumin.

## **SURGICAL TECHNIQUE**

All patients were given systemic antibiotics (tablet ciprofloxacin 500mg b.d.) on the preoperative day. On the day of surgery pupils were dilated adequately using instillation of 0.8% tropicamide and 5% /10% phenylephrine eye drops every 10 minutes, one hour before surgery. To sustain the pupil dilatation the anti-prostaglandin eye drops such as flubiprofen should be instilled three times one day before surgery and half hourly for two hours immediately before surgery.



## **SURGICAL STEPS OF MANUAL SMALL INCISION CATARACT SURGERY**

1. The eye to be operated is painted, draped and prepared for surgery under aseptic precautions.
2. Local anesthesia is given using 2% xylocaine mixed with 1500 units of hyaluronidase.
3. Universal wire speculum applied.
4. Superior rectus (bridle) suture is passed to fix the eye in downward gaze.
5. A small fornix based conjunctival flap is made, and sclera is exposed.
6. Haemostasis is achieved by applying gentle and just adequate wet field cautery.
7. A self sealing sclera-corneal tunnel incision is made.
8. Side-port entry is made with the help of 1.5mm valvular corneal incision at 9<sup>o</sup> clock position.
9. Anterior capsulotomy by continuous curvilinear capsulorrhexis of adequate size is done.
10. Hydrodissection is done to separate cortico-nuclear mass from the posterior capsule.
11. Depending on the degree of mydriasis the pupil was stretched mechanically or sphincterotomies were done, depending on the operating surgeon's discretion.
12. Synechiolysis was done if required
13. Nucleus was delivered.
14. Cortical matter was removed by irrigation and aspiration.
15. In case of a posterior capsule tear, the integrity of the capsular bag was assessed to place the intraocular lens.
16. In case of vitreous loss, manual anterior vitrectomy was done.
17. If there were no complications, posterior chamber intraocular lens was placed in the capsular bag.

18. The viscoelastic was cleared from the anterior chamber.
19. Subconjunctival gentamycin and dexamathasone 0.5cc was given at the end of the procedure.
20. Pad and bandage applied.

Postoperatively all the patients received a course of topical antibiotic and steroid eye drops one hourly .systemic antibiotic was given for 5 days postoperatively.

## **OBSERVATION AND RESULTS**

---

## RESULTS AND OBSERVATION

**Table 1**

### AGE DISTRIBUTION IN PATIENTS WITH PSEUDOEXFOLIATION SYNDROME

(n = 100)

AGE	NUMBER OF PATIENTS	PERCENTAGE
50 – 59 years	26	26
60 – 70 years	50	50
71 – 80 years	24	24
Total	100	100




Figure 2: Age distribution in patients with Pseudoexfoliation Syndrome

In the present study, there were 26 (26.00 %) patients of age group 50 – 59 years, 50 (50%) patients of age group 60 – 70 years and 24 (24.00 %) of age group 71. The average age of patients was 65.83 years and about 73 (73.00 %) of patients were above 60 yrs of age.

**Table 2:**

**SEX DISTRIBUTION IN PSEUDOEXFOLIATION SYNDROME**

(n = 100)

SEX	NUMBER OF PATENTS	PERCENTAGE
Male	63	63.00
Female	37	37.00
Total	100	100




Figure 3: Sex Distribution in Pseudoexfoliation Syndrome

In the present study, there were 63 (63.00%) were males and 37 (37.00%) were females.

**Table 3:**

**LATERALITY IN EYES WITH PSEUDOEXFOLIATION SYNDROME**

(n = 100)

LATERALITY	NUMBER OF PATIENTS	PERCENTAGE
Unilateral	50	50
Bilateral	50	50
Total	100	100





Figure 4: Laterality in patients with Pseudoexfoliation Syndrome

In the present study there were 50 (50%) of patients had clinical bilateral involvement of Pseudoexfoliation syndrome and 50 (50%) had unilateral involvement.

**Table 4:**

**IRIS CHARACTERISTICS IN PSEUDOEXFOLIATION SYNDROME**

CHARACTERISTICS	PRESENT	ABSENT
Iris Surfcae	40	60
Pupillary Margin	80	20
Moth Eaten appearance	48	52
Iridodonesis	1	99
Atrophy	10	90
Posterior Syncechiae	12	88



**Figure 5: Iris Characteristics in Pseudoexfoliation Syndrome**


In the present study, 80% of patients had Pseudoexfoliation material on the pupillary margin, 40 (40%) on the iris surface, 46 (46.00 %) have Moth Eaten Appearance, 10 (10 %) have Iris Atrophy, 1 (0.33 %) have Iridodonesis and 4 of the patients had posterior synechiae.

**Table 5:**

**ANGLE CONFIGURATION IN PSEUDOEXFOLIATION SYNDROME**

(n = 100)

ANGLE CONFIGURATION	NUMBER OF PATIENTS	PERCENTAGE
Open angles	90	90
Narrow angles	10	10
Total	100	100



**Figure 6: Angle Configuration in Pseudoexfoliation Syndrome**

In the present study, of the 100 patients with Pseudoexfoliation syndrome, 90(90%) of patients had open angles and 10(10%) patients had narrow angles.



**Table 6:**

**INTRAOCULAR PRESSURE IN PSEUDOEXFOLIATION SYNDROME**

(n = 100)

IOP	NUMBER OF PATIENTS	PERCENTAGE
$\leq 21$ mm Hg	70	70
22 – 30 mm Hg	23	23
31 – 40 mm Hg	4	4
$\geq 41$ mm Hg	3	3
Total	100	100




Figure 7: Intraocular Pressure in Pseudoexfoliation Syndrome

In this study, out of 100 patients in the present study group, 70 (70%), 23 (23.00%), 4 (4.00%) and 3 (3.00%) of the patients had intraocular pressure  $\leq 21$  mm of Hg, between 22 mm of Hg to 30 mm of Hg, between 30 mm of Hg to 40 mm of Hg and  $\geq 41$  mm of Hg respectively.

**Table 7:**

**PUPILLARY DILATATION IN PSEUDOEXFOLIATION SYNDROME**

(n = 100)

PUPILLARY DILATATION	NUMBER OF PATIENTS	PERCENTAGE
Sufficient (> 6 mm)	53	53.00
Insufficient (<6 mm)	47	47.00
Total	100	100




Figure 8: Pupillary dilatation in Pseudoexfoliation Syndrome


In the present study, 53 (53.00 %) of patients had sufficient mydriasis, and 47 (47.00 %) of the patients had insufficient mydriasis in eyes with Pseudoexfoliation syndrome.

**Table 8:**

**ZONES OF PSEUDOEXFOLIATION ON LENS SURFACE**

(n = 100)

ZONE	NUMBER OF PATIENTS	PERCENTAGE
Peripheral & Central Zone	40	40
Only Peripheral Zone	60	60
Only Central Zone	0	0
Total	100	100



**Figure 9: Zones of Pseudoexfoliative Lens Material**

In the present study, 60 (60 %) had peripheral zone, 40 (40 %) had both peripheral zone and central zone and none of them had only central zone.

**Table 9:**

**TYPE OF CATARACT IN PSEUDOEXFOLIATION SYNDROME**

(n = 100)

TYPE OF CATARACT	NUMBER OF PATIENTS	PERCENTAGE
SMC	30	30
NS I	3	3.00
NS II	27	27
NS III	40	40
TOTAL	100	100




Figure 10: Type of Cataract in Pseudoexfoliation Syndrome

In the present study, 30 (30%) of the patients had Mature Cataract, 3(3.00%) had NS grade1 cataract, 27(27.00%) of the patients had NS grade2 cataract and 40(40%) of the patients had NS grade 3 cataract.

**Table 10:**

**INTRAOPERATIVE COMPLICATIONS IN PSEUDOEXFOLIATION SYNDROME**

(n = 100)

COMPLICATIONS	NUMBER OF PATIENTS	PERCENTAGE
OCCURRED	16	16.00
NOT OCCURRED	84	84.00
TOTAL	100	100





Figure 11: Intraoperative Complications in Pseudoexfoliation Syndrome

In the present study, 16 (16.00 %) of the patients had intraoperative complication while 84 (84.00 %) did not.

**Table 11:**

**INDIVIDUAL TYPES OF INTRAOPERATIVE COMPLICATIONS IN PATIENTS WITH PSEUDOEXFOLIATION SYNDROME**

COMPLICATION	NUMBER OF PATIENTS	PERCENTAGE
Zonular dehiscence	10	10.00
Posterior capsular rent	10	10.00
Vitreous loss	9	9.00



**Figure 12: Individual Types of Intra-operative Complications in Pseudoexfoliation Syndrome**

As shown in Table 11, 10 (10.00 %) of patients had Zonular dehiscence, 10(10.00 %) of the patients had Posterior Capsular Rent and 9 (9.00 %) of the patients had Vitreous loss.

**Table 12:**

**SURGICAL OUTCOME IN PATIENTS WITH PSEUDOEXFOLIATION SYNDROME**

(n = 100)

OUTCOME	NUMBER OF PATIENTS	PERCENTAGE
Intraocular Lens Implantation	90	90
Aphakic Plane lens extraction	10	10
Total	100	100





Figure 13: Surgical Outcome in Patients with Pseudoexfoliation Syndrome

In the present study of Pseudoexfoliation syndrome, 90 (90 %) of the patients were implanted with intraocular lens after employment of various surgical modifications like Sphincterectomy, Sphincterotomy, manual Anterior Vitrectomy. 10 (10 %) of the patients were left aphakic due to the above mentioned complications

**Table 13:**

**INTRAOPERATIVE COMPLICATIONS VERSUS PUPILLARY  
DILATATION**

	INSUFFICIENT MYDRIASIS	ADEQUATE MYDRIASIS
Posterior Capsular Rent	6	4
Vitreous loss	5	4
Zonular dehiscence	6	4



**Figure 14: Intraoperative Complications versus Pupillary Dilatation**


In the present study, Out of the total 100 patients, 16 (16.00 %) patients who had one or the other or all of the complications, 62 (62.00%) of them had insufficient mydriasis and 38 (38.00 %) of the patients had adequate mydriasis.



**Table 14:**

**SURGICAL MODIFICATIONS IN PATIENTS WITH  
PSEUDOEXFOLIATION SYNDROME**

MODIFICATION	NUMBER OF PATIENTS
Sphincterectomy	24
Synechiolysis	13
Both	10



**Figure 15: Surgical Modification in Patients with Pseudoexfoliative Syndrome**

In the present study, 13 (13.00 %) underwent Synechiolysis, 24 (24.00 %) patients underwent sphincterotomy and 10 (10 %) patients underwent both sphincterotomy and Synechiolysis.

## **DISCUSSION**

---

## **DISCUSSION**

This study consisted of 100 eyes of 100 patients with Pseudoexfoliation syndrome who underwent manual small incision cataract surgery in R L J HOSPITAL AND RESEARCH CENTER, TAMAKA, KOLAR attached to Sri Devaraj Urs Medical college.

As shown in table 1, there were 27 (27 %) patients of age group 50 – 59 years, 50 (50%) patients of age group 60 – 70 years and 23 (23 %) of age group 71 – 80 %. The average age of patients was 65.83 years and about 73 (73.00 %) of patients were above 60 yrs of age. The prevalence of Pseudoexfoliation syndrome increases with age. Pseudoexfoliation syndrome usually occurs between 60 to 80 yrs, the average age being 70 yrs. In this study, 73.00 % of the patients are in the age group of 60 – 80 years which is in concurrence with the mentioned studies.

As shown in table 2, 63 (63.00%) were males and 37 (37.00%) were females. Studies regarding the sex distribution of Pseudoexfoliation syndrome are conflicting. Women have predominated in some series while other studies have found equal or greater prevalence in men.

As shown in table 3, 50 (50% ) of patients had clinical bilateral involvement of Pseudoexfoliation syndrome and 50 (50%) had unilateral involvement. A review of literature comparing the frequency of monocular versus binocular involvement in various series is not conclusive. Many series have reported bilateral involvement to be more common with ratios as high as 3:1 while other studies have reported unilateral involvement to predominate again with ratios as high as 3:1.

**Hammer, Schlotzer- Schrehardt, Naumann**<sup>55</sup> in 2001 carried out an ultrastructural study of the contralateral eye in 5 pairs of donor eyes with unilateral Pseudoexfoliation syndrome. They showed “ultrastructural” alterations in anterior segment tissues of all the eyes. They concluded that basically Pseudoexfoliation syndrome is a bilateral disease with clinically marked asymmetric manifestations. The reasons for this marked asymmetry remain unknown. Clinically unilateral involvement is often a precursor to bilateral involvement within 5- 10 yrs after diagnosis.

As shown in table 4, 80% of patients had Pseudoexfoliation material on the pupillary margin, 40 (40%) on the iris surface, 46 (46%) have Moth Eaten Appearance, 10 ( 10 % ) have Iris Atrophy, 1 ( 1 % ) have Iridodonesis, 13(13%) had posterior synechiae in this study group. This is in concurrence with the study by **Ritch Schlotzer. Scherhardt**<sup>56</sup> (2001) stated that deposits of Pseudoexfoliation material on the iris sphincter and pupillary margin are seen in 84% patients. Thus next to the lens Pseudoexfoliation material, the most prominent and consistent clinical finding is the Pseudoexfoliation material at the pupillary border.

In the present study, 43 ( 43 %) of patients had average trabecular pigmentation, and 33 (33 %) of patients had ‘moderate trabecular pigmentation’ and 24 (24 %) of patients had ‘heavy pigmentation’. None of the patients had ‘Absent Or Faint Pigmentation’.

The extent of trabecular pigmentation has been correlated to the degree of Increased intraocular pressure.

In this present study, 23 (23 %) of the patients had Pseudoexfoliation material in the angle and the same was absent in 77 (77 %) of the patients. In the present study is very close by to the study conducted by **Sunde** (1956).

**Sunde** (1956) found flakes of Pseudoexfoliation material in the angle in 18.75% of the patients with pseudoexfoliative glaucoma. But **Tarkkanen** (1962) found Pseudoexfoliation material in the angle on gonioscopic examination in 46% of glaucomatous eyes and 50% of non glaucomatous eyes. Thus the two conditions can occur simultaneously or separately and the relation between them is not clear.

**Schlotzer-Scherhardt et al** (1992) found a significant correlation between the extent of Pseudoexfoliation material in the angle and Pseudoexfoliation syndrome.

As shown in Table 5, of the 100 patients, 90(90% ) of patients had open angles and 10(10% ) patients had narrow angles. Out of 10 narrow angles, 4 patient had narrow angle due to anterior subluxation of the lens, 3 patient had narrow angles due to synechial angle closure and 3 patient had narrow angles due to appositional angle closure. The present study is in concurrence with the following studies.

In studies of patients with Pseudoexfoliation syndrome, occludable angles were noted in 9-18% of patients (**Bruce shields** 1999, fourth edition).

**Ritch, Schlotzer- Scherhardt** (2001) noted 23% of patients with Pseudoexfoliation syndrome and glaucoma to have grade 2 or narrow angles.

**Wishartetal**<sup>57</sup> (1985) noted 32% of patients in their study to have narrow angles.

As shown in Table 6, out of 100 patients in the present study group, 70 (70 %), 23 (23 %), 4 (4%) and 3 (3%) of the patients had intraocular pressure  $\leq$  21 mm of Hg, between 22 mm of Hg to 30 mm of Hg, between 30 mm of Hg to 40 mm of Hg and  $\geq$  41 mm of Hg respectively.

Out of these patients, 20 (20 %) had Primary Open angle glaucoma and 10 (10%) had Secondary angle closure glaucoma.

In patients with pseudoexfoliation syndrome, 20% have glaucoma and increased IOP at the time of diagnosis. Patients who have pseudoexfoliation syndrome but not glaucoma should be considered vulnerable to glaucoma, because 15% of such patients develop increased IOP within 10 years. This underscores the need for careful follow-up in patients who have pseudoexfoliation syndrome. Pseudoexfoliation syndrome accounts for 15-20% of cases of open angle glaucoma. The present study is in concurrence with this finding.

As shown in Table 7, 53 (53 %) of patients had sufficient mydriasis, and 47 (47 %) of the patients had insufficient mydriasis. This is in concurrence with the following studies.

**Freyler H, Radax U** (1990) noted pupillary dilatation less than 4 mm in 19 of 32 patients with pseudoexfoliation syndrome who underwent cataract surgery.

**Asano N, Schlotze – Scherhardt, Naumann** (1996) attributed poor mydriasis in Pseudoexfoliation syndrome to degenerative changes of sphincter and dilator muscle tissues and apparent involvement of the muscle cells in Pseudoexfoliation material fiber formation.

**Repo L.P. et al** (1996) found degenerative changes in both the stromal tissue and in the muscular layer of iris and regarded this as one of the causes for miosis.

**Alfaite et al** (1996) noticed significant insufficient mydriasis (p value < 0.001) in their study of 31 patients with Pseudoexfoliation syndrome.

Reduction of stromal elasticity by accumulation of pseudoexfoliation material may also play a role in poor mydriasis.

**Stanila A** (1996) also noted an increased incidence of insufficient pupil dilatation in the 10 patients with Pseudoexfoliation syndrome undergoing cataract surgery in their study.

**Avramides S, Trainanidis P, Sakkias G** (1997) in their study of 84 patients with Pseudoexfoliation syndrome who underwent ECCE, noted that 61.90% of them had pupillary dilatation less than 5 mm.

In the present study, 20 % had pigment dispersion after mydriasis. **Prince, A.M., Ritch R** <sup>59</sup> (1986) reported that anterior chamber melanin dispersion after mydriasis may be seen as a whorl like pattern of pigment particles on iris sphincter and peripheral iris.

**Ritch R, Schlotze – Scherhardt** (2000) reported pigment dispersion in the anterior chamber after mydriasis to be common and profuse in Pseudoexfoliation syndrome. Pigment dispersion after mydriasis is one of the suspicious sign to meticulously look out for Pseudoexfoliation syndrome in preclinical stages.

As shown in Table 8, 60 (60 %) had peripheral zone, 40 (40 %) had both peripheral zone and central zone and none of them had only central zone. The peripheral zone of pseudoexfoliation material is a consistent finding and the central zone is not always apparent (**M. Bruce Shields**). **Tarkkanen** (1962) found the central zone absent in 18% of cases in his study while **Ritch, Schlotzer – Scherhardt** (2001) found it absent in 20 – 60% of their cases.

As shown in Table 9, 30 (30%) of the patients had Mature Cataract and 70 (70 %) of them had Immature cataract. All of them, i.e. 100 %, had Nuclear Cataracts. Cortical Cataract was present along with advanced nuclear cataract and none of the patients had isolated cortical cataract.

**Seland et al**<sup>60</sup> (1982) have reported a higher incidence of nuclear cataract in eyes with pseudoexfoliation syndrome with fewer cortical cataracts. **Hietanen J. et al** have also reported nuclear cataract to be the predominant type of cataract in Pseudoexfoliation syndrome.

**Ritch R, Schlotze – Scherhardt** (2001) have also reported an increased incidence of nuclear cataract in Pseudoexfoliation syndrome.

In the present study, 10 (10%) of patients had phacodonesis. **Futa R. Furnyoshi**<sup>61</sup> (1989) reported an 8.4% incidence, while **Moreno J., Duch S., Harara J** (1993) reported a 10.6% incidence of phacodonesis. 3 (3 %) of the patients had iridodonesis. This is because the iris in Pseudoexfoliation syndrome is more rigid due to vascular compromise and various other changes like deposition of Pseudoexfoliation material, Atrophy, Loss of iris stroma – Moth Eaten Appearance.



As shown in Table 10, 27 (27 %) of the patients had intraoperative complication while 73 (73 %) did not. As shown in Table 11, 10 (10 %) of patients had Zonular dehiscence, 10 (10 %) of the patients had Posterior Capsular Rent and 9 (9%) of the patients had Vitreous loss. As shown in Table 12, 90 (90%) of the patients were implanted with intraocular lens after employment of various surgical modifications like with iris hooks, Sphincterotomy, Synecholysis, manual Anterior Vitrectomy. 10 (10 %) of the patients were left aphakic due to the above mentioned complications.

**Scrolloli et al**<sup>62</sup> (1998) have found that PEX patients were five times more likely to develop intraoperative complications during cataract surgery compared to patients without the condition.

**Schönherr U et al** (1989) found a statistically significant increase in intraoperative and postoperative complication in eyes with Pseudoexfoliation syndrome in their study of 436 patients.

**Freyler H, Radax U** (1990) found 26 of their 36 patients with pseudoexfoliation syndrome undergoing ECCE to have intraoperative complication—Zonular dehiscence, Posterior capsular rent and Vitreous loss.

Various studies in eyes with Pseudoexfoliation syndrome have quoted the incidence of Zonular dehiscence to be 17.90%. **Hovding G** (1998), 13.1% by **Avramides S** (1997) and 14.8% by **Lumme P, Laatikanan** (1993).

**Alfaiete et al** (1996) in their study of 31 patients found zonular dehiscence to be more common in eyes with Pseudoexfoliation syndrome but this was not statistically significant when compared to eyes without Pseudoexfoliation syndrome.

**Stanila** (1996) also reported an increased incidence of Posterior capsular rent and Vitreous loss in their study of 10 eyes with Pseudoexfoliation syndrome undergoing ECCE.

**Kuchle et al** (2000) found 6.9% of their 11 patients to have intraoperative complication namely – zonular dehiscence and vitreous loss. Zonular fragility in Pseudoexfoliation syndrome increases the risk of lens dislocation, zonular dehiscence and vitreous loss up to 10 times (Ritch R, 2001).

**Lumme P, Laatikainen L** (1993) found the incidence of vitreous loss to be fourfold more in eyes with Pseudoexfoliation syndrome and Posterior capsular rent to be 10 fold higher in eyes with Pseudoexfoliation syndrome.

**Avramides S., Travamides P, Sakkias G** (1997) found the incidence of Posterior capsular rent and vitreous loss to be 10.4% and 7.14% respectively in this study of 84 patients with Pseudoexfoliation syndrome undergoing cataract surgery.

The incidence of vitreous loss in eyes with Pseudoexfoliation syndrome undergoing cataract surgery has been reported by various authors as 11.9% by **Kuchle** (1989) and 6.7% by **Junemann A, Mart us P, et al** (1997).

**Naumann G.O., Kucle M. Schonher U** (1978) in their study of 72 eyes with Pseudoexfoliation syndrome found a seven fold increase for vitreous loss in eyes with Pseudoexfoliation syndrome as compared to those without Pseudoexfoliation syndrome. They also noted the incidence of posterior capsular rent to be 4.2% in eyes with Pseudoexfoliation syndrome and 2.8% without Pseudoexfoliation syndrome.

As shown in Table 13, out of 10 (10 %) of the patients with posterior capsular rent, 6 (60.00%) of them had insufficient mydriasis while 4 (40.00 %) of them had adequate mydriasis. Out of 9 (9 %) of the patients who had Vitreous loss, 5 (55.45%) of them had insufficient mydriasis while 4 (44.55 %) of them had adequate mydriasis. Of the 10 (10 %) of the patients with Zonular dehiscence, 6 (60.00 %) of them had insufficient mydriasis while 4 (40.00 %) of them had adequate mydriasis. Out of the total 100 patients, 16 (16 %) patients who had one or the other or all of the above complications, 10 (61.75. %) of them had insufficient mydriasis and 6 (38.25 %) of the patients had adequate mydriasis. In the present study, out of total 8 patients, 62.5 % of them had inadequate mydriasis.

This correlates with the other studies conducted by **Freyler H., Radax U** (1990), **Asano N. et al** (1996), **Repo L.P. et al** (1996), **Stanilla A.** (1996) and **Avramides S et al** (1997).

As shown in Table 14, 13 (13 %) underwent Synechiolysis, 23 (23 %) patients underwent sphincterotomy and 10 (10 %) patients underwent both sphincterotomy and Synechiolysis.

**Alfaite et al** (1996) in their study of 31 eyes of Pseudoexfoliation syndrome undergoing ECCE noted a statistically significant increase ( $p$  value  $< 0.01$ ) in the need to perform sphincterotomies.

**Kuchle et al** (2000) noted 3.4% of their 76 patients to require surgical Synechiolysis and/or mechanical dilatation of pupil intraoperatively.

**Vickie Lee and Anthony Maloof** (2002) studied extensively on small pupils and their management during cataract surgery. They advocated that small

pupils could be enlarged by prosthetic and non – prosthetic methods. Non – prosthetic techniques include visco-mydriasis, manual iris stretching and iris micro- sphincterotomies. Prosthetic techniques include iris hooks and use of pupil expansion devices.

In the present study, out of the 16 patients who had one or the other or all of the above mentioned intraoperative complications, 6 of them had Senile Mature Cataract, 6 of them had Senile Hypermature Cataract while the remainder 4 had Senile Immature Cataract. The patient with hyper mature cataract had Zonular dehiscence, Posterior capsular rent, Vitreous loss, Corneal edema by the end of the surgery and excessive lens mobility while performing capsulotomy. This is in concurrence with the study mentioned below.

**Bayramlar et al** (2007) conducted a retrospective study in 225 eyes of 187 patients of which 99 eyes had Pseudoexfoliation syndrome. Pre-operative data collected were – Cataract maturity level, Best corrected Visual acuity and Intraoperative posterior capsule complications. In this study, he interpreted that in MSICS, Pseudoexfoliation syndrome has an increased intraoperative posterior capsule complication rate that increases as the level of cataract maturity increases. Thereby, it is advisable to operate early on cataracts in patients with Pseudoexfoliation syndrome to have better results and prevent the compromised Zonular and posterior capsule changes.

## **CONCLUSION**

---

## CONCLUSIONS

The following conclusions were drawn from the study.

Patients with pseudoexfoliation syndrome and cataract posted for manual small incision cataract surgery, have to be carefully looked for zonular weakness, insufficient mydriasis, IOP, subluxation or dislocation of cataractous lens because these preoperative factors have bearing on the intraoperative complications.

Inadequate mydriasis is one of the major pre operative complications in eyes with Pseudoexfoliation syndrome which has a bearing on the intr operative complications like posterior capsular rent and vitreous loss.

Adequate surgical modifications such as Sphincterotomy and/or synechiolysis, pupil stretching, use of iris hooks in these eyes with inadequate mydriasis reduce the intra operative complications. These pupil enlargement procedures are advocated during cataract surgery.

All though cataract surgery in Pseudoexfoliation syndrome is challenging, if the surgeon is aware of the condition pre operatively and pays meticulous attention to the surgical technique during manual small incision cataract surgery, the intr aoperative complications can be managed and a good outcome can be expected.

Manual small incision cataract surgery is safe in eyes with pseudoexfoliation syndrome.

## **SUMMARY**

---

## SUMMARY

In the present study titled “THE STUDY OF INTRAOPERATIVE COMPLICATIONS OF MANUAL SMALL INCISION CATARACT SURGERY IN EYES WITH PSEUDOEXFOLIATION SYNDROME”, 100 eyes of 100 patients with both Cataract and Pseudoexfoliation syndrome, attending R L JALLAPA hospital and research center, attached to Sri Devraj Urs medical college, Tamaka, Kolar were included. The study included patients of average age group of these patients was 65.83 years with preponderance of males with equal incidence of unilateral and bilateral involvement.

In the present study, majority of the patients had Pseudoexfoliation material on the pupillary margin with 100% patients having different grades of trabecular pigmentation with 90% of the patients having open angles, 70% patients had IOP within normal range, 20 % of the patients had open angle glaucoma and 10% having secondary angle closure glaucoma. 46.67 % of the patients had insufficient mydriasis. 100 % of the patients had either peripheral zone or central zone or both of the pseudoexfoliation material on the anterior surface of the lens suggesting Pseudoexfoliation syndrome. 30 % of patients had cataract maturity level more than nuclear sclerosis grade 4.

In the present study, 16 % of the patients had either Zonular dehiscence, Posterior capsular rent, Vitreous loss, Cornealedema, Excessive lens mobility during capsulotomy and combination of the above. Majority of these patients with intraoperative complications had insufficient mydriasis and higher cataract maturity level. The surgical modifications like Sphincterotomy, Synechiolysis,



Manual anterior vitrectomy, use of CTR can improve the outcome of the surgery and give better visual quality to the patients with Pseudoexfoliation syndrome.

## **BIBLIOGRAPHY**

---

## BIBLIOGRAPHY

1. **Lindberg JG.** Kliniska undersökningar över depigmentering av pupillarranden och genomylsbarkefav iris vid fall av alderstarr samit inormala ogon hos gamla personer. Doctoral thesis, Diss Helsingfors, 1917.
2. **Vogt A.** Ein neues Spaltlampenbild des Pupillargebietes: Hellblauer Pupilearsaumfillz mit Nautchenbildunz aus der Lisenvorderkapsel. Klin Monatsbl Augenheilkd 75:1, 1925.
3. **Busacca A.** Struktur und Bedeutung der Hautschennieder-Schlaze in der vorderen und hinteren Augendammer. Graefes Arch Clin Exp Ophthalmol 1927; 119:13335.
4. **Dvorak-Theobald G.** Pseudoexfoliation of the lens capsule: Relation to "true" exfoliation of the lens capsule as reported in the literature and role in the production of glaucoma capsulocuticulare. Trans Am Ophthalmol Soc 1953;51: 387.
5. **Ashton N, Shakib M, Collyer R, Blach R.** Electron microscopic study of pseudoexfoliation of the lens capsule. I. Lens capsule and zonular fibers. Invest Ophthalmol 1965; 4:141.
6. **Bertelsen TI, Drablos PA, Flood PR.** The so-called senile exfoliation (pseudoexfoliation) of the anterior lens capsule, a product of the lens epithelium. Acta Ophthalmol 1964; 42: 1096.
7. **Eagle RC Jr., Font RL, Fine BS.** The basement membrane exfoliation syndrome. Arch Ophthalmol 1979; 97:510
8. **Sunde OA.** Senile exfoliation of the anterior lens capsule. Acta Ophthalmol, 1956; 45:1.

9. **Layden WE, Shaffer RN.** Exfoliation syndrome. *Am J Ophthalmol* 1974; 78:835.
10. **Tarkkanen A, Forsius H, eds.** Exfoliation syndrome. *Acta Ophthalmol* 1988;66(suppl. 184):1.
11. **Christopher P et al.** “Diagnosis and Management of Pseudoexfoliation glaucoma” *Eye net* 2006.
12. **Sood N.N.** “Prevalence of Pseudoexfoliation of the lens capsule in India”. *Acta Ophthalmol.*1968; 46: 211-214.
13. **Lamba P.A. and Giridhar A.** “Pseudoexfoliation syndrome (prevalence based on random survey hospital data)” *Indian Journal of Ophthalmology* 1984; 32: 169 – 173.
14. **Aravind H et al.** “Pseudoexfoliation in South India” *British Journal of Ophthalmology* 2003; 87 (11): 1321 – 1323.
15. **Tarkkanen A.** “Pseudoexfoliation”. *Acta Ophthalmol Suppl.* 1962; 71: 1 – 98.
16. **Kelvin Y.C. Lee et al.** “Association of LOXL1 polymorphisms with Pseudoexfoliation in the Chinese” *Molecular Vision* 2009; 15: 1120 – 1126.
17. **R.R. Allingham et al.** “Pseudoexfoliation syndrome in Icelandic families” *British Journal of Ophthalmology*, 2001; 85: 702 – 707.
18. **M. Bruce Shield’s** Text book of Glaucoma, 5th edition, Lippincott Williams & Wilkins, Philadelphia.
19. **Deepak Gupta.** Glaucoma Diagnosis and Management, Chapter 16 “Pseudoexfoliation syndrome”, Lippincott Williams & Wilkins © 2005, Philadelphia, USA.

20. **Jack J Kanski.** Chapter 13 ‘Glaucoma’, “clinical Ophthalmology” 6<sup>th</sup> edition, Printed in UK; Copyright © 2007 by Elsevier’s health sciences rights department, 397-399.
21. **L C Datta.** section 66 “pseudoexfoliation syndrome”, Modern Ophthalmology 1st edition., printed in India; copyright © 1994 by Jaypee brother’s medical publishers pvt.ltd.,463.
22. **Mizuno K and Muroi S.** “Cycloscopy of pseudoexfoliation.” American journal of ophthalmology 1979; 87: 513.
23. **Yanoff and Duker.** Ophthalmology 3rd edition, Chapter 10.13, “Pseudoexfoliative glaucoma.” Copyright © 2008 Mosby. An imprint of Elsevier.
24. **William E. Layden,** Becker Schaffer’s The Glaucoma, Chapter 8 Exfoliation syndrome, Mosby Publication, © 1989, page 997 – 1013.
25. **Schlotzer-Schrehardt et al.** “pseudoexfoliation syndrome –ocular manifestation of a systemic disorder?” Archives of ophthalmology, 1992;110(12) 1752-1756.
26. **Schumacher S. et al.**“Pseudoexfoliation syndrome and Aneurysms of Abdominal aorta”, The Lancet, 2009; 357: 359 – 360.
27. **Harnish J P, Barrach H J, Hassel J R, et al.** “Identification of a Basement membrane proteoglycan in Pseudoexfoliation material”. Graefe’s Archcin Experimental Ophthalmology, 1981; 215 – 273.
28. **Garner, A and Alexander, RA.** Pseudoexfoliative disease: histochemical evidence of an affinity with zonular fibres, British Journal of Ophthalmology 1984: 68; 574.

29. **Streeten BW et al.** "Pseudoexfoliative fibrillopathy in the conjunctiva: a relation to elastic fibres and elastosis" – *Ophthalmology* 1987; 94:1439 – 1449.
30. **Schlotzer – Schrehardt U, Naumann G O, Kuchle M.** Pseudoexfoliationsyndrome for the comprehensive ophthalmologist. Intraocular and systemic manifestations. *Ophthalmology*, 1998; 105: 951-68.
31. **Repo L.P. Naucharinen et al.** "Pseudoexfoliation syndrome with poorly dilating pupil: a light and electron microscopic study of the sphincter area". *Graefes Archcin Experimental Ophthalmology* 1996; 234 (3):171 – 176.
32. **Tsukahara and Matsuo T.** "Secondary glaucoma accompanied with primary familial amyloidosis", *Ophthalmologica* 1977: 175: 250.
33. **Mizuno K, Hara S, Ishiguru S and Takei Y.** "Acid phosphatase in eyes with Pseudoexfoliation", *American journal of Ophthalmology*, 1980; 89: 482.
34. **Baba H.** "Histochemical and polarization – Optical investigation for glycosoaminoglycans in Pseudoexfoliation syndrome", *Graefe's Archcin Experimental Ophthalmology*, 1983; 221: 106.
35. **Davanger M.** "A note on the Pseudoexfoliation fibrils" *Acta Ophthal* 1978;56: .114.
36. **Davanger M.** "Studies on the Pseudoexfoliation material" *Graefe's Archcin Experimental Ophthalmology*, 1978; 208: 65.

37. **Skuta G. L., Parrish R. K. et al.** “Zonular dialysis during Extra capsular cataract surgery in pseudoexfoliation syndrome.” *Arch Ophthalmol* 1987; 105(5): 632-634.
38. **Naumann G. O., Kuchle M. Schonherr U.** “Pseudoexfoliation syndrome as a risk factor for vitreous loss in extra-capsular cataract extraction”. *Fortschr ophthalmol* 1989; 86:543-545.
39. **Wang L, Yamasita R and Hammura S.** “corneal endothelial cell changes and aqueous flare intensity in eyes with pseudoexfoliation syndrome.” ***Ophthalmologica*** 1999;213: 387-391.
40. **Kuchle M, Naumann.H et al.** “pseudoexfoliation syndrome and secondary cataract.” *British journal of ophthalmology* 1997; 81:862-866.
41. **Lumme P, Lattikainen L.** “Exfoliation syndrome and cataract extraction.” *American journal of ophthalmology* 1993; 116(1): 51-55.
42. **Moreno M.J., Duch S and Lajara.** “pseudoexfoliation syndrome :Clinical factors related to capsular rupture in cataract surgery.”. *Acta Ophthalmol (Copenh)* 1993; 71: 181 – 184.
43. **Freyler H and Radax U.** “pseudoexfoliation syndrome –a risk factor in modern cataract surgery?.” *Klin Monatsbl Augenheilkd* 1994;205: 275-279.
44. **Stanila A.** “The exfoliation syndrome. The risk factor in Extra-capsular surgery of the crystalline lens.” *Ophthalmologica* 1996;40(4); 373-376.
45. **Kuchle et al.** “Anterior chamber depth and complications during cataract surgery in eyes with pseudoexfoliation syndrome.” *American journal of ophthalmology* 2000;129: 281-285.

46. **Bayramlar H, Hepsen F, Yilmaz.H.** “Mature cataracts increase risk of capsular complications in manual small incision cataract surgery in pseudoexfoliative eyes”, *Can j ophthalmology* 2007, February;42(1);46-50.
47. **Albert Galand , Michael Kuchle , Etienne Hachet.** “multiple surgical challenges of pseudoexfoliation” Symposium during 21st congress of the ESCRS 2004.
48. **Vickie Lee and Anthony Maloof.** “Clinical practice –Cataract surgery in pseudoexfoliative syndrome” *Comprehensive Ophthalmology update* 2002,3(1).
49. **Howard Fine.** “Pseudoexfoliation: A double Challenge”, *ASCRS Eye World*, 2008; 10:10.
50. **Roger F Steinert.** Chapter 1 “The pathology of Cataract”, . *Cataract surgery* 2nd edition. , 7.
51. **Norman S Jaffe et al.** “Cataract surgery and its complication”, 6th edition, Mosby – A Harcourt Health Sciences Company., 71, 301.
52. **Aravind Eye hospital.** Chapter 5 “Complications of Manual Small Incision Cataract surgery and their management”, © 2000 by Aravind Publications, Aravind Eye Hospitals., 35 – 42.
53. **Devgan Uday.** “Cataract Surgery – Pseudoexfoliation can create challenges in cataract surgery”. *OSN Supersite* – 2008.
54. **Cionni R.J., Osher R.H.** “Endocapsular Ring approach to the subluxated cataractous lens.”, *Journal of Cataract and Refractive Surgery*, 1995; 21: 245 – 249.



55. **Hammer, Schlotzer- Schrehardt, Naumann.** “Unilateral or Asymmetric Pseudoexfoliation syndrome? An electron microscopic study”., *Klin Monatsbl Augenheilkd* 2001;216(6): 388-392.
56. **Ritch Schlotzer, Scherhardt et al.** “Unilateral or Asymmetric Pseudoexfoliation syndrome, An Ultra structural Study”, *Achieves of Ophthalmology*, 2001; 119: 1023 – 1031.
57. **Wishart PK, Spathe GL and Poryzees EM.** “Anterior Chamber angle in the Exfoliation Syndrome”, *British Journal of Ophthalmology*, 1985; 69:103.
58. **Asano N, Schlotze – Scherhardt, Naumann GO.** “A histo-pathological study of iris changes in Pseudoexfoliation”, *Ophthalmology*, 1996; 102: 1279 – 1290.
59. **Prince, A.M., Ritch R and Streeten BW.** “Preclinical diagnosis of Pseudoexfoliation”, *Archives of Ophthalmology*, 1987: 105: 1076.
60. **Seland GH and Chylack LT Jr.** “Cataracts in the Exfoliation Syndrome (Fibrilliopathia epitheliocapsularis)”., *Transophthalmol Soc U.K.*,1982; 102: 375.
61. **Futa R, Furnyoshi N, Shimizu T.** “Clinical features of capsular glaucoma in comparision with primary open angle glaucoma in Japan”. *Acta Ophthal* 1992;70: 214-219.
62. **Scorolli L et al.** “Pseudoexfoliative syndrome – A cohort study on intraoperative complications in cataract surgery”, *International Journal of Ophthalmology*, 1998; 212 (4): 278 – 280.

63. **David J.Apple et al.** “Intraocular lenses”, Evolution, designs, complications and pathology printed in USA copyrights © 1989, Williams and Wilkins; 120, 144, 386.
64. **Menkhaus S et al.** “Pseudoexfoliation syndrome and intraoperative complications in Cataract surgery”, *Klin Monatsbl Augenheilkd*, 2000; 216 (6): 388 – 392.
65. **Miller K.M., Kener G.T. Jr.** “Stretch pupiloplasty for small pupil phacoemulsification”, *American Journal of Ophthalmology*, 1994, 117; 107 – 108.

# **ANNEXURES**

**ANNEXURE I: PROFORMA FOR THE STUDY OF INTRAOPERATIVE COMPLICATIONS OF MANUAL SMALL INCISION CATARACT SURGERY IN EYES WITH PSEUDOEXFOLIATION.**

CASE NUMBER:

NAME:

AGE:      SEX:      I.P. NO./O.P. NO.:

DATE OF ADMISSION:      DATE OF DISCHARGE:

FINAL DI AGNOSIS:

PRESENTING COMPLAINTS:

HISTORY OF PRESENTING COMPLAINTS:

GENERAL PHYSICAL EXAMI NATION:

OCULAR EXAMINATION:

<b>EXAMINATION</b>	<b>RIGHT EYE</b>	<b>LEFT EYE</b>
VISUAL ACUITY		
DISTANT VISION <ul style="list-style-type: none"> <li>- Uncorrected</li> <li>- Pin Hole</li> <li>- Corrected</li> </ul>		
NEAR VISION		
ANTERIOR SEGMENT EVALUATION		
ADNEXA		
CONJUNCTIVA		
SCLERA		
CORNEA		
ANTERIOR CHAMBER <ul style="list-style-type: none"> <li>- AC Depth</li> <li>- Pigment Dispersion</li> </ul>		

<b>EXAMINATION</b>	<b>RIGHT EYE</b>	<b>LEFT EYE</b>
<b>IRIS</b> <ul style="list-style-type: none"> <li>- Colour</li> <li>- Pattern</li> <li>- Moth eaten appearance</li> <li>- Posterior synechiae</li> <li>- Others</li> </ul>		
<b>PUPIL</b> <ul style="list-style-type: none"> <li>- Size</li> <li>- Shape</li> <li>- Reaction</li> <li>- Direct</li> <li>- Indirect</li> <li>- PEX on Pupillary margin</li> </ul>		
<b>LENS</b> Type of Cataract Zones of PEM Phacodonesis Subluxation Dislocation Others		
<b>FUNDUS</b>		
Direct Ophthalmoscopy		
Indirect Ophthalmoscopy		
<b>INTRA- OCULAR PRESSUR</b>		
By Schiottz Tonometry		
<b>LACRIMAL PATENCY TEST</b>		

<b>EXAMINATION</b>	<b>RIGHT EYE</b>	<b>LEFT EYE</b>
<b>GONIOSCOPY</b>		
Angle ( by Schaffer's Classification)		
Trabecular pigmentation		
Pseudoexfoliative material		
Sampolesi's Line		
Others		
<b>POST-MYDRIASIS</b>		
Pupil size		
Pigment Dispersion		
<b>EXTRA-OCULAR MUSCLE MOVEMENTS</b>		
<b>VISUAL FIELD</b>		
By Confrontation Method		

A SCAN ESTIMATION OF POWER OF IOL

PRE-OPERATIVE PREPARATION:

OPERATIVE NOTES:

DATE:

PROCEDURE:

SURGEON:

INTRA-OPERATIVE-COMPLICATIONS:

Zonular Dialysis

Posterior Capsular rupture

Need for Sphincterotomy

Need for Synechiolysis

Vitreous Loss

Others

**ANNEXURE II: MASTER CHART**

Sl. No.	IP. No.	Age	Sex	UL/BL	Iris characteristics						Pupil					Lens					
					Surface	PM	Moth	PS	Atrophy	I/D	Pre-dilatation			Post-dilatation		Type of cataract	Zones		PD	SL/DL	ZD
											Size	Shape	Reaction	Size	PID		pz	pz+cz			
1	450823	60	F	UL	-	+	-	-	+	-	3	r	N	7	+	NS GR3	+	-	-	-	-
2	451154	58	F	UL	-	+	+	-	-	-	3	r	N	5	-	NSGR2+SC	-	+	-	-	-
3	451591	70	M	UL	-	+	+	-	-	+	4	r	SR	5	+	SHMC	+	-	+	+	+
4	222968	80	M	UL	+	+	+	+	-	-	2	r	SR	6	+	NSGR2+SC	-	+	+	-	+
5	450226	78	M	BL	+	-	-	-	-	-	3	r	N	5	+	NSGR3	-	+	-	-	+
6	433241	70	F	UL	-	-	-	-	+	-	3	irre	N	7	-	SMC	-	+	-	-	-
7	432744	67	M	UL	-	-	+	-	-	-	3	r	N	6	+	NSGR2	-	+	-	-	-
8	453201	55	M	BL	+	+	+	-	-	-	3	r	N	7	+	NSGR2	+	-	-	-	+
9	453200	72	F	BL	-	+	-	-	-	-	2	r	SR	6	-	NSGR3	-	+	-	-	-
10	453210	50	M	UL	-	+	+	-	-	-	2	r	SR	5	-	NSGR2	-	+	-	-	-
11	448778	70	F	BL	-	+	-	-	+	-	3	r	N	7	-	NSGR3	-	+	-	-	-
12	453758	80	M	UL	+	+	-	-	-	-	3	r	N	7	+	SMC	+	-	-	-	-
13	454778	65	M	BL	+	+	+	-	-	-	3	r	N	7	+	NSGR3	+	-	-	-	-
14	454177	70	M	BL	-	+	-	-	-	-	3	r	N	7	-	NSGR1	-	+	+	+	+
15	439469	70	M	UL	-	+	-	-	-	-	3	r	N	7	-	NSGR3	-	+	-	-	-
16	454784	80	M	UL	-	+	-	-	-	-	3	r	N	6	-	NSGR2	+	-	-	-	-
17	454778	65	M	UL	+	+	-	-	-	-	2	r	SR	8	+	SMC	-	-	-	-	-
18	454811	60	F	UL	+	+	-	-	-	-	3	r	N	6	+	NSGR3	-	-	-	-	-
19	455573	70	M	BL	+	+	-	-	-	-	3	r	N	8	+	SMC	+	+	-	-	-
20	321801	68	F	BL	-	+	-	-	-	-	3	r	N	7	-	SMC	+	-	-	-	-
21	454747	60	F	BL	-	-	+	-	-	-	2	r	SR	7	-	NSGR3	+	-	-	-	-
22	455704	58	M	BL	-	-	+	-	-	-	3	r	N	6	-	NSGR3	-	-	-	-	-

23	456240	75	M	BL	-	+	+	-	-	+	2	r	SR	5	+	SMC	+	+	-	-	-
24	457443	55	F	BL	-	+	-	+	-	+	2	irre	SR	5	+	SMC	-	-	-	-	-
25	454577	70	M	UL	-	+	+	+	-	+	3	irre	SR	5	-	SMC	-	-	-	-	-
26	457798	80	M	BL	+	+	+	+	-	-	2	irre	N	6	-	NSGR3	+	+	-	-	-
27	455632	50	M	UL	+	-	-	-	-	-	3	r	N	9	-	NSGR3	-	+	-	-	-
28	460550	64	F	BL	+	-	+	-	-	-	2	r	SR	7	-	NSGR3	-	+	-	-	-
29	460202	50	F	UL	-	+	+	-	-	-	3	r	N	9	-	NSGR2	-	+	-	-	-
30	458269	55	M	BL	-	+	-	-	-	-	3	r	N	9	+	NSGR3	+	+	-	-	+
31	454784	55	M	UL	+	+	-	-	-	-	2	r	N	6	-	SMC	+	-	-	-	+
32	454817	60	M	BL	+	+	-	-	-	-	3	r	N	7	-	SMC	+	-	-	-	-
33	460459	57	M	UL	-	+	-	-	-	-	2	r	N	6	-	NS1	+	-	-	-	-
34	460573	61	M	BL	-	+	-	-	-	-	3	r	N	7	+	NS2	-	+	-	-	-
35	462224	64	F	BL	+	+	-	-	-	+	3	r	N	7	+	NS1	-	+	-	-	-
36	462229	65	F	UL	-	-	-	-	+	+	3	r	N	7	+	NS2	-	+	-	-	-
37	456507	66	M	UL	-	-	-	-	-	-	3	r	N	8	+	NS3	+	-	-	-	-
38	462227	51	F	BL	+	-	-	-	+	-	4	r	N	7	+	NS3	+	-	-	-	-
39	463304	68	M	BL	+	+	+	+	+	-	4	r	N	8	+	NS2	+	-	+	-	-
40	431910	71	M	BL	-	+	+	+	+	-	2	r	N	8	-	NS3	-	-	+	-	-
41	431187	58	M	UL	-	+	+	+	+	-	2	r	N	8	-	NS2	-	-	+	-	-
42	464748	66	M	UL	+	+	+	-	-	-	3	r	N	8	+	NS3	+	+	-	-	-
43	465163	69	M	UL	-	+	-	-	-	+	2	irre	SR	8	+	SMC	+	-	-	-	-
44	464844	61	F	BL	-	+	-	-	-	+	3	irre	SR	7	-	SMC	-	-	-	-	+
45	465162	54	F	BL	+	-	-	+	-	-	2	r	N	6	-	SMC	-	-	-	-	+
46	465637	64	M	UL	-	-	-	+	-	-	2	r	N	6	-	SMC	+	-	-	-	+
47	465622	55	M	UL	+	-	-	-	+	-	2	r	N	8	+	NS3	+	+	-	-	-
48	465619	68	M	UL	-	-	-	-	+	-	3	r	N	8	+	NS3	-	+	-	-	-
49	465639	58	M	UL	-	+	+	-	-	-	3	r	N	7	+	NS3	-	-	-	-	-



50	466797	64	F	BL	+	+	-	-	-	-	3	r	N	7	_	NS2	+	-	-	-	-
51	467241	57	F	BL	_	+	-	-	-	-	4	r	N	6	_	NS2	+	-	-	-	-
52	459669	67	M	BL	-	+	+	-	-	+	4	r	N	6	_	NS3	+	+	-	-	-
53	467248	54	M	UL	-	+	+	-	+	+	3	r	N	9	+	SMC	+	+	-	-	-
54	467236	66	F	UL	+	-	-	+	+	+	3	r	N	8	+	SMC	+	+	-	-	-
55	467242	68	F	UL	+	-	-	+	-	-	3	irre	SR	8	+	NS2	-	-	+	-	-
56	467218	55	F	BL	_	-	-	+	-	-	2	irre	SR	7	_	NS3	+	-	+	-	+
57	467318	65	M	BL	-	-	+	-	-	-	3	r	N	7	_	NS2	+	-	+	-	+
58	466619	54	M	UL	-	+	+	-	-	-	2	r	N	8	+	NS3	-	-	-	-	+
59	465639	69	M	UL	+	+	+	-	+	-	2	r	N	8	_	SMC	+	-	-	-	-
60	466797	55	F	UL	-	+	-	+	+	-	3	r	N	9	_	SMC	+	+	-	-	-
61	467241	61	F	BL	-	+	-	+	-	-	3	r	N	9	+	NS3	+	+	-	-	-
62	468693	52	M	BL	+	+	+	+	-	-	3	r	N	8	_	NS3	+	-	-	-	-
63	468707	64	M	BL	+	+	+	-	-	-	2	r	N	7	_	NS2	-	-	-	-	-
64	468700	52	M	UL	+	+	+	-	-	-	2	r	N	5	+	NS2	-	-	-	-	-
65	396248	63	M	UL	-	+	-	-	+	-	3	r	N	5	+	SMC	+	-	-	-	-
66	467979	53	F	BL	-	-	-	-	+	-	3	r	N	6	+	SMC	+	-	-	-	-
67	467981	69	F	BL	-	-	-	-	+	-	3	r	N	7	_	NS3	+	-	-	+	-
68	470204	55	M	UL	-	-	+	-	-	+	2	r	N	7	+	NS3	-	-	-	+	-
69	468489	62	M	UL	-	+	+	-	-	+	2	r	N	6	_	NS2	-	-	-	+	-
70	470200	59	M	UL	+	+	+	-	-	-	2	r	N	8	_	NS3	+	+	-	-	-
71	468712	70	M	UL	+	+	-	+	-	-	3	r	N	8	_	NS2	+	-	+	-	-
72	468704	68	F	BL	+	-	-	+	-	-	3	r	N	9	_	SMC	+	-	+	-	+
73	470196	64	M	UL	+	-	-	+	-	-	3	r	N	7	_	SMC	-	-	+	-	+
74	470201	65	F	UL	-	-	+	-	-	-	2	r	N	6	+	NS3	-	-	-	-	-
75	469810	69	M	BL	-	+	-	-	-	-	2	r	N	6	+	NS2	+	+	-	-	-
76	470281	61	F	UL	-	+	+	-	-	-	2	r	N	6	_	NS2	+	-	-	-	-

77	470315	59	M	BL	-	+	-	-	-	-	3	r	N	8	-	NS3	+	-	-	-	-
78	471493	76	F	UL		+	+	-	-	-	3	r	N	7	-	NS2	-	+	-	-	
79	471499	72	M	BL	-	+	--	-	-	-	4	r	N	9	+	NS3	+	-	-	-	-
80	471780	78	F	UL	-	+	--	-	+	-	4	r	N	9	-	NS2	+	+	-	-	-
81	471775	66	F	BL	+	-	-	+	+	-	3	r	N	7	-	NS3	-	-	-	-	+
82	479346	77	M	UL	-	-	+	+	+	+	3	r	N	7	+	SMC	-	+	-	-	-
83	471782	68	M	BL	+	-	+	-	-	+	2	irre	SR	8	+	SMC	+	+	-	-	-
84	471776	62	M	BL	+	-	+	-	-	+	2	irre	SR	8	-	NS3	+	+	-	-	-
85	464935	74	M	UL	-	+	-	-	+	-	3	r	SR	6	-	NS2	+	+	-	-	+
86	451117	79	F	BL	-	+	-	+	+	-	3	r	N	6	+	NS2	+	+	-	-	-
87	473490	68	M	BL	+	+	-	+	-	-	2	r	N	8	+	NS3	-	+	-	-	-
88	473478	64	M	BL	+	-	+	+	-	-	2	r	SR	7	-	NS3	+	+	-	-	+
89	473477	65	F	BL	+	-	+	+	+	+	3	r	SR	8	+	NS2	+	-	-	-	-
90	473484	80	F	BL	-	-	-	-	+	+	3	r	N	8	-	SMC	+	-	-	-	-
91	462146	79	M	BL	-	-	+	-	-	-	3	r	N	7	-	SMC	+	-	-	-	-
92	477444	76	F	UL	-	+	-	-	-	-	4	r	N	6	-	NS3	+	+	+	-	+
93	475125	77	M	BL	-	+	-	-	+	+	4	r	N	6	+	NS3	+	-	+	-	-
94	475113	74	F	BL	+	+	+	-	-	+	2	r	N	8	+	NS3	+	-	+	-	-
95	475128	76	M	UL	+	+	+	-	+	-	3	r	N	8	+	NS2	+	-	-	+	-
96	475136	79	F	BL	-	+	-	+	-	-	2	irre	N	8	+	NS3	+	-	-	+	+
97	476676	74	M	BL	-	+	+	+	-	+	3	r	SR	9	-	SMC	+	-	-	-	+
98	476701	76	F	UL	+	-	-	+	-	-	3	irre	SR	6	+	SMC	+	-	-	-	-
99	476684	75	M	BL	+	-	+	-	-	-	2	r	N	7	+	SMC	-	-	-	-	-
100	473477	66	F	BL	-	+	-	-	-	+	2	irre	SR	8	-	NS2	-	-	-	-	-

Sl. No.	IP. No.	OP (mm Hg)	Gonioscopy				Surgery							
			Config	Tmp	PAS	PxM	Executed	Outcome	Modification		Complications			
									Shp.	Syn.	ZD	PCR.	VL	Others
1	450823	20.6	Open	2+	-	+	MSICS	Iol	-	-	-	-	-	-
2	451154	23.1	Open	4+	+	+	MSICS	Iol	-	-	-	-	-	-
3	451591	24.5	Narrow	4+	-	-	MSICS	Aphakic	+	-	-	+	+	CO+ELM
4	222968	12.6	Open	3+	-	+	MSICS	Iol	-	+	+	-	-	-
5	450226	18.0	Open	3+	-	+	MSICS	Aphakic	+	-	-	-	-	-
6	433241	17.3	Open	2+	-	-	MSICS	Iol	-	-	-	-	-	-
7	432744	43.4	Open	4+	-	+	MSICS	Iol	-	-	-	-	-	-
8	453200	11.2	Open	2+	-	-	MSICS	Iol	+	-	+	-	-	-
9	453210	23.0	Narrow	2+	-	-	MSICS	Iol	-	-	-	-	-	-
10	448778	18.0	Open	4+	-	-	MSICS	Iol	-	-	-	-	-	-
11	453758	14.6	Open	2+	-	-	MSICS	Iol	-	-	-	-	-	-
12	448999	20.6	Open	3+	-	-	MSICS	Iol	-	-	-	-	-	-
13	439469	17.3	Open	2+	-	-	MSICS	Iol	-	-	-	-	-	-
14	454177	20.6	Open	2+	-	-	MSICS	Iol	-	-	-	-	-	-
15	454778	31.8	Open	3+	-	-	MSICS	Iol	-	-	-	-	-	-
16	454784	15.9	Open	2+	-	-	MSICS	Iol	-	-	-	-	-	-
17	454788	17.0	Open	2+	-	-	MSICS+Trab	Aphakic	+	-	-	+	+	ELM
18	454811	17.3	Open	3+	-	-	MSICS+Trab	Iol	-	-	-	-	-	-
19	455573	20.6	Open	2+	-	-	MSICS	Iol	-	-	-	-	-	-
20	321801	17.3	Open	2+	-	-	MSICS	Iol	-	-	-	-	-	-
21	454747	14.6	Open	4+	+	+	MSICS	Iol	-	-	-	-	-	-
22	455704	20.6	Open	3+	-	-	MSICS	Iol	-	-	-	-	-	-

23	456240	19.6	Open	3+	-	-	MSICS	Iol	+	+	-	-	-	-
24	457443	21.0	Open	2+	-	-	MSICS	Iol	+	+	-	-	-	-
25	454577	30.0	Open	4+	-	-	MSICS	Iol	+	+	-	-	-	-
26	457798	28.0	Open	3+	+	+	MSICS	Iol	-	+	-	+	-	-
27	454632	19.6	Open	2+	-	-	MSICS	Iol	-	-	-	-	-	-
28	458614	27.0	Open	3+	-	-	MSICS	Iol	-	-	-	-	-	-
29	459297	23.0	Open	3+	-	-	MSICS	Iol	-	-	-	-	-	-
30	459450	19.6	Open	4+	-	-	MSICS	Iol	-	-	-	-	-	-
31	458269	43.0	Narrow	2+	-	-	MSICS	Aphakia	+	+	+	-	+	CO
32	460202	13.0	Open	2+	-	+	MSICS	Iol	-	-	-	-	-	-
33	460550	16.0	Open	2+	-	-	MSICS	Iol	-	-	-	-	-	-
34	460459	33.0	Narrow	3+	-	-	MSICS	Iol	-	-	-	-	-	-
35	460573	36.0	Narrow	4+	+	-	MSICS	Iol	-	-	-	-	-	-
36	462224	39.0	Narrow	3+	+	-	MSICS	Iol	-	-	-	-	-	-
37	462229	23.0	Open	3+	-	-	MSICS	Iol	+	-	-	-	-	-
38	456507	16.0	Open	2+	-	+	MSICS	Iol	-	+	+	-	-	-
39	462227	13.0	Open	4+	-	+	MSICS	Iol	-	-	-	-	-	-
40	463315	24.0	Open	4+	-	-	MSICS	Iol	-	-	-	-	-	-
41	463304	15.0	Open	3+	-	-	MSICS	Iol	-	-	-	-	-	-
42	431910	12.0	Open	3+	+	-	MSICS	Iol	-	-	-	-	-	-
43	431187	24.0	Open	2+	+	-	MSICS	Iol	-	-	-	-	-	-
44	464748	15.0	Open	4+	-	-	MSICS	Iol	-	-	-	-	-	-
45	465163	13.0	Open	4+	-	-	MSICS	Iol	-	+	-	-	-	-
46	464844	21.0	Open	3+	-	+	MSICS	Iol	+	-	-	-	-	-
47	465162	11.0	Open	3+	-	-	MSICS	Iol	-	-	-	-	-	-
48	465637	28.0	Open	2+	-	-	MSICS	Iol	-	-	-	-	-	-
49	465622	13.0	Open	3+	-	-	MSICS	Iol	-	-	-	-	-	-


50	465619	23.0	Open	3+	-	-	MSICS	Iol	+	-	-	-	-	-
51	465639	14.0	Open	4+	-	-	MSICS+Trab	Iol	+	+	-	-	+	-
52	466797	13.0	Open	4+	-	-	MSICS	Iol	-	-	-	-	-	-
53	467241	25.0	Narrow	3+	-	-	MSICS	Iol	-	-	-	-	-	-
54	459669	12.0	Open	3+	-	-	MSICS	Aphakia	+	+	-	+	+	CO
55	467235	15.0	Open	3+	+	+	MSICS	Iol	-	-	-	-	-	-
56	467235	25.0	Open	2+	-	-	MSICS	Iol	-	-	-	-	-	-
57	467233	12.0	Open	3+	-	-	MSICS	Iol	-	-	-	-	-	-
58	467248	22.0	Open	4+	-	-	MSICS	Iol	-	-	+	-	-	-
59	467236	16.0	Open	2+	+	-	MSICS	Iol	-	-	-	-	-	-
60	467242	12.0	Open	3+	+	-	MSICS	Iol	-	+	-	+	+	-
61	467218	13.0	Open	4+	-	-	MSICS	Iol	+	-	-	-	-	-
62	467218	22.0	Open	3+	-	-	MSICS	Iol	+	-	-	-	-	-
63	467318	14.0	Open	3+	-	+	MSICS	Iol	-	-	+	-	-	-
64	467979	12.0	Open	3+	-	+	MSICS	Iol	-	-	-	-	-	-
65	467981	24.0	Open	2+	-	-	MSICS	Iol	-	-	-	-	-	-
66	396248	17.0	Open	3+	+	-	MSICS	Iol	-	-	-	-	+	-
67	468700	16.0	Open	4+	-	-	MSICS	Iol	-	+	-	-	-	-
68	468707	22.0	Open	3+	-	-	MSICS	Iol	-	-	+	-	-	-
69	468893	12.0	Open	2+	-	+	MSICS	Iol	-	-	-	-	-	-
70	470204	22.0	Open	2+	+	-	MSICS	Iol	+	-	-	-	-	-
71	468469	16.0	Open	3+	-	-	MSICS	Aphakia	+	+	-	+	+	CO
72	470200	14.0	Open	4+	-	+	MSICS+Trab	Iol	-	-	-	--	-	-
73	468712	24.0	Open	3+	-	+	MSICS	Iol	-	-	-	-	-	-
74	468704	12.0	Open	3+	-	-	MSICS	Iol	-	-	-	-	-	-
75	470196	15.0	Open	2+	-	-	MSICS	Iol	-	-	-	-	-	-
76	470201	24.0	Open	4+	+	-	MSICS	Iol	-	-	-	-	-	-

77	469810	11.0	Open	2+	-	-	MSICS	Iol	-	-	+	-	-	-
78	470281	16.0	Open	3+	-	-	MSICS	Iol	-	-	-	-	-	-
79	470315	25.0	Open	4+	-	-	MSICS	Iol	-	-	-	-	-	-
80	471493	14.0	Open	3+	-	-	MSICS	Iol	-	-	-	-	-	-
81	471499	14.0	Open	3+	+	-	MSICS	Iol	-	-	-	-	-	-
82	471780	12.0	Open	4+	+	-	MSICS	Aphakia	+	+	-	+	-	-
83	471775	15.0	Open	3+	-	-	MSICS	Iol	-	-	-	-	-	-
84	479346	19.0	Open	2+	-	-	MSICS	Iol	-	-	-	-	-	-
85	471782	15.0	Open	3+	-	-	MSICS+Trab	Iol	+	-	-	-	-	-
86	471776	14.0	Open	3+	-	+	MSICS	Aphakia	+	+	+	+	-	-
87	464935	15.0	Open	2+	-	-	MSICS	Iol	-	-	-	-	-	-
88	451117	12.0	Open	4+	-	-	MSICS	Iol	-	-	-	-	-	-
89	473490	22.0	Open	3+	-	-	MSICS	Iol	+	-	-	-	-	-
90	473490	16.0	Open	3+	+	-	MSICS	Aphakia	+	-	-	+	-	-
91	473477	14.0	Open	2+	+	+	MSICS	Iol	-	-	-	-	-	-
92	473484	14.0	Open	3+	-	+	MSICS	Iol	-	-	-	-	-	-
93	462374	28.0	Narrow	4+	-	-	MSICS	Iol	-	-	-	-	-	-
94	462146	16.0	Open	3+	-	-	MSICS	Iol	-	-	-	-	-	-
95	477444	14.0	Open	2+	-	-	MSICS+Trab	Iol	+	-	+	-	-	-
96	475125	13.0	Open	3+	+	-	MSICS	Iol	-	-	-	-	-	-
97	475138	14.0	Open	4+	-	-	MSICS	Iol	-	-	-	-	-	-
98	475113	15.0	Open	3+	-	-	MSICS	Iol	-	-	-	-	-	-
99	475128	12.0	Open	3+	-	-	MSICS	Aphakia	+	-	-	+	+	CO
100	476701	16.0	Open	3+	-	+	MSICS	Iol	-	-	-	-	-	-


### ANNEXURE III: KEY TO MASTER CHART

BL » Bilateral	PCR » Posterior capsular rent
CO » Corneal edema	PD » Phacodonesis
Config. » Configuration	PID » Pigment dispersion
cz » Central zone	PM » Pupillary margin
D/L » Dislocation	PS » Posterior synechiae
ELM » Excessive lens mobility	pz » Peripheral zone
F » Female	R » Regular
GR » Grade	S/L » Subluxation
I/D » Iridodonesis	SC » Subcapsular cataract
IOL » Intraocular lens	SHMC » Senile Hypermature Cataract
IOP » Intraocular pressure	Sl. No. » Serial number
IP. No. » Inpatient number	SMC » Senile mature cataract
irre » Irregular	Sph. » Sphincterotomy
M » Male	SR » Sluggishly reactive
mm Hg » Millimeter of Mercury	Syn. » Synechiolysis
Moth » Moth eaten appearance	Tmp » Trabecular pigmentation
MSICS » Manual Small Incision Cataract Surgery	Trab » Trabeculectomy
N » Normal	UL » Unilateral
NS » Nuclear sclerosis	VL » Vitreous loss
PAS » Peripheral anterior synechiae	yrs » Years
	ZD » Zonular dehiscence

**ANNEXURE IV: PHOTOGRAPHS**



Photograph 1: Pseudoexfoliation material at the pupillary margin and anterior surface of lens capsule with undilated pupil.




Photograph 2: Zone of Pseudoexfoliation material on the anterior capsule of lens after pupillary dilatation.







Photograph 3: Moth Eaten Appearance in Pseudoexfoliation syndrome



Photograph 4: Histopathology of Pseudoexfoliation material on the anterior lens capsule.



Photograph 5: Sphincterotomy cut at 10 o'clock position. Seen postoperatively.



Photograph 6: Capsular Tension Ring.