



Effects of Barley Semolina on Physiological And Biochemical Parameters in Chronic Kidney Disease Patients

KEYWORDS

Barley semolina, nutritional status, renal functions

* Dr. Madhavi Reddy

Clinical Nutritionist, Department of medicine,
Sri Devaraj Urs Medical College, Tamaka, Kolar,
Karnataka-563 101

Raja Reddy P

Assistant professor, Department of Physiology,
Sri Devaraj Urs Medical College, Tamaka, Kolar,
Karnataka-563 101

Raghavendra Prasad BN

Professor, department of Medicine, Sri Devaraj Urs
Medical College, Tamaka, Kolar, Karnataka - 563 101

Mahesh V

Assistant professor in Community Medicine, Sri Devaraj
Urs Medical College, Tamaka, Kolar - 563101.

ABSTRACT

Scanty data is available in the dietary use of barley in chronic kidney disease patients. **Objectives:** To study the effects of barley semolina on physiological and biochemical parameters in chronic kidney disease patients. **Methods:** Forty-nine patients of stage three chronic kidney disease were selected to evaluate the effects of barley semolina. Baseline patient's data including anthropometric measurements, hemogram, serum electrolytes, lipid profile and renal function were noted. All participants were then advised to consume barley in their daily breakfast at 10% of daily calorie requirement for one month and 20% in the second month. At the end of both first and second month the participants were evaluated for change in baseline data. **Results:** A significant improvement in the nutritional status was observed at both 10% and 20% of nutritional intervention with barley. The total cholesterol and triglycerides were reduced ($p < 0.001$) and HDL cholesterol increased with both interventions. Sodium was significantly increased ($p < 0.001$) with the 20% intervention only. Potassium decreased with both the interventions ($p < 0.001$). Hemoglobin values were improved at 10% and 20% intervention levels. **Conclusion:** Introduction of barley in the diet of stage 3 chronic kidney disease patients has significantly improved the nutritional status and renal functions. Lowered total cholesterol, triglycerides and increased HDL cholesterol were also found. This is a hypothesis generating study and further work needs to be done in this area on a large scale.

INTRODUCTION

The role of nutritional support in kidney disease is to prevent or reserve associated malnourished status, minimize the adverse effects of substances that are inadequately excreted and favorably affect the progressions and outcome of kidney disease. Careful dietary management may make it possible to stabilize the progression of chronic kidney disease and avoid or postpone dialysis¹. Conservative management of chronic kidney disease (CKD) has included variety of dietary manipulations and supplementation of dietary fiber to reduce adverse symptoms is a novel approach².

Barley is one of the cereal grains and is a staple food in most countries of Middle East. In other western countries it is grown for animal feed and for malt extract. Barley in India is used as human food and barley flour is by and large mixed with wheat or gram flour for preparing chapattis and also in making bread³. Barley is source of dietary fiber, b-glucan and antioxidants⁴. Barley is a good diuretic among cereals, barley water made after cooking cereal is very beneficial in all urinary and kidney disease conditions. It is rich in amino acid L-tryptophan, vitamin E, minerals such as calcium, copper, iodine, magnesium and potassium³.

Barley intake has been shown to have beneficial effects on lipid metabolism. Clinical data continue to demonstrate that the consumption of barley products is effective in lowering total and LDL-cholesterol^{4,5}. Chronic kidney disease patients are advised low sodium diet to avoid volume over loads and hemodynamic instability and low potassium diet to avoid the potential complications of hyperkalemia. The chromium content of barley has shown to improve glucose tolerance in glucose intolerant people and could be of benefit in dietary management of diabetes mellitus⁵. Limited studies are available on the effects of barley on chronic kidney disease. In view of this, the present study was undertaken to study the

health benefits of barley cereal in chronic kidney disease.

Objectives

To study the effects of barley semolina on nutritional status, lipid profile and kidney functions in patients with chronic kidney disease.

MATERIALS AND METHODS

The study was conducted in RL Jalappa Hospital and research Centre, Kolar, Karnataka. A total of 92 patients fitting into the selection criteria (CKD stage 3, MDRD calculation) attending medicine outpatient department from June 2009 to July 2010 were enrolled. Informed consent was obtained from the patients willing to participate in the study. Anthropometric measurements have been used as a marker of nutritional status and body composition. Measurements like body weight, height, skin fold thickness and mid arm muscle circumference were recorded. Body mass index calculated and also biochemical tests conducted as per standard lab procedure and recorded.

Cleaned and conditioned barley is milled to make semolina. In the beginning of the study Uppuma and porridge prepared of barley was given to ten patients to test the acceptability in terms of appearance, texture, taste, color and flavor⁶. The recorded scores were within acceptable range. Patients and their attendants were demonstrated the use of semolina in preparing Uppuma and porridge for the breakfast. For each patient the daily calorie requirement was calculated. 10% of it was substituted with barley to be consumed as breakfast. Patient wise daily pouches of 10% barley semolina was calculated and distributed for one month. The patients were followed up weekly enquiring about their barley consumption through telephone contact. At the end of the one month, anthropometric measurements and biochemical tests were performed and findings recorded. Barley consumption was

increased to 20% of the daily calorie requirements for the second month and similar follow up was done. Out of 92 patients enrolled for the study 49 patients completed the two months follow-up. Other 43 patients were dropped out in between due to their noncompliance in follow-up. At both the clinic visits, demography data, personal history, physical activity and medications used were determined with a standard questionnaire. Medications were validated by examination of prescriptions or pills brought to the physician.

The data obtained was analyzed using SPSS software version 16, change variables were defined as follow-up values minus baseline values. Specific mean differences between follow-up and baseline visits were employed by paired t-test to test the significant difference between the samples.

RESULTS

Forty-nine chronic kidney disease patients participated in the study, 21(42.85%) male and 28(57.15%) were female (fig.1). Introduction of barley cereal in the diet of stage 3 CKD patients improved the kidney function, the mean values of serum urea, creatinine and uric acid were not significant both at 10% and 20% of barley intervention (table 1). Total cholesterol and triglycerides were reduced significantly ($P < 0.001$) both at 10% and 20% barley intervention. At the same time HDL cholesterol increased at 10% and 20% barley intervention ($p < 0.001$). No change of sodium at 10% intervention but at 20% intervention it was increased. Potassium decreased at both 10% and 20% intervention table 2. Significant increase in nutritional status were observed in weight, MAMC and body mass index at 10% and 20% intervention, but there was no improvement in SFT at 10% intervention, slight difference were observed at 20% intervention (table 3&4). While total protein, albumin and hemoglobin were not significant both at 10% and 20% barley intervention (table 1&2).

DISCUSSION

There is an improvement in the nutritional status of the patients in the study as the body weight and BMI were elevated significantly at 10% and 20% of dietary intervention. Whereas serum albumin, total protein and hemoglobin levels were not significant at 10% and 20% of dietary intervention. A study from National Health and Nutritional Examination Survey (NHANES) revealed CKD patients with improved nutritional status have better survival rate⁷. Most people with diabetes and CKD require sodium restriction in diet to have optimum blood pressure control and to prevent fluid retention⁸. In the present study no change in sodium levels was observed at 10% barley diet intervention, while at 20% intervention it was raised. Diuretic property of the barley may be the reason for improvement in the kidney function in the study⁹. In this regard, there is a need to optimize renal bone disease management through prescription of phosphate binding medications and vitamin D derivatives¹⁰.

The present study showed a statistical significant ($p < 0.001$) decrease serum cholesterol and triglycerides and at the same time a statistically significant ($p < 0.001$) increased HDL levels. Lupton JR et al concluded that addition of barley bran flour or barley oil enhances the cholesterol lowering effect¹¹. The ability of barley B-glucan soluble fiber to lower serum cholesterol is a combination of factors and mechanisms. Suggested mechanisms of cholesterol reduction include, delayed intestinal absorption of glucose and lipids and inhibition of absorption and reabsorption of cholesterol and bile acids accompanied by increased excretion of bile acids^{12,13,14}. The reduced absorption may be caused by the high viscosity of b-glucan solutions, which increases the viscosity of the intestinal contents^{15,16,17}. Other factors may also be important, such as the fermentation of b-glucan in the colon, resulting in production of short-chain fatty acids, which impede cholesterol biosynthesis¹³.

CONCLUSION

The effects of barley semolina in the diet of stage 3 CKD pa-

tients had significant difference for total cholesterol, triglycerides and HDL-c. At the same time significant difference was observed for nutritional status. However, further research in this field needs to be accompanied in CKD patients, to substantiate the health benefits of barley.

Table.1 Mean values of biochemical parameters at 10% barley intervention

Parameters	Base line Mean \pm SD	10% invention Mean \pm SD	p-value
Urea(mg/dl)	60.35 \pm 12.72	56.12 \pm 1.45	0.001*
Creatinine(mg/dl)	4.26 \pm 1.21	3.68 \pm 1.05	0.001*
Uric acid(mg/dl)	4.87 \pm 1.06	4.34 \pm 0.86	0.437
Total protein(g)	5.47 \pm 0.93	5.97 \pm 0.95	0.543
Albumin(g)	4.19 \pm 0.74	4.73 \pm 0.87	0.001*
Hemoglobin(g)	10.9 \pm 2.02	11.15 \pm 2.05	0.655
Cholesterol(mg/dl)	204.33 \pm 35.57	196.02 \pm 32.7	0.001*
Triglycerides(mg/dl)	190.35 \pm 65.96	182.65 \pm 59.61	0.001*
HDL-c (mg/dl)	33.39 \pm 5.26	34.06 \pm 4.84	0.001*
Sodium(mEq/l)	141.43 \pm 13.92	141.43 \pm 13.92	0.234
Potassium(mEq/l)	5.28 \pm 0.85	4.85 \pm 0.71	0.001*

* $P < 0.001$ considered as significant

Table.2 Mean values of biochemical parameters at 20% barley intervention

Parameters	Baseline Mean \pm SD	10% invention Mean \pm SD	p-value
Urea(mg/dl)	60.35 \pm 12.72	54.9 \pm 12.82	0.001*
Creatinine(mg/dl)	4.26 \pm 1.21	2.06 \pm 0.69	0.001*
Uric acid(mg/dl)	4.87 \pm 1.06	3.37 \pm 0.8	0.243
Total protein(g)	5.47 \pm 0.93	7.39 \pm 0.62	0.453
Albumin(g)	4.19 \pm 0.74	5.11 \pm 0.7	0.001*
Hemoglobin(g)	10.9 \pm 2.02	21.33 \pm 1.92	0.876
Cholesterol(mg/dl)	204.33 \pm 35.57	190.53 \pm 20.77	0.001*
Triglycerides(mg/dl)	190.35 \pm 65.96	187.59 \pm 154.59	0.001*
HDL-c(mg/dl)	33.39 \pm 5.26	35.04 \pm 4.33	0.001*
Sodium(mEq/l)	141.43 \pm 13.92	147.55 \pm 11.03	0.457
Potassium(mEq/l)	5.28 \pm 0.85	4.05 \pm 0.53	0.001*

* $P < 0.001$ considered as significant

Table 3. Mean values of nutritional status at 10% barley intervention

Parameters	Baseline Mean \pm SD	10% intervention Mean \pm SD	p-value
Weight(kg)	60.57 \pm 10.31	61.57 \pm 10.09	0.001*
Height(cm)	162.92 \pm 8.08	162.92 \pm 8.08	0.231
SFT(mm)	16.94 \pm 2.96	16.94 \pm 2.96	0.061
MAMC(cm)	19.22 \pm 2.66	19.44 \pm 2.66	0.072
BMI(kg/m ²)	22.72 \pm 2.84	23.11 \pm 2.73	0.001*

* $P < 0.001$ considered as significant

Table 4. Mean values of nutritional status at 20% barley intervention

Parameters	Baseline Mean±SD	20% intervention Mean±SD	p-value
Weight(kg)	60.57±10.31	62.47±8.97	0.001*
Height(cm)	162.92±8.08	162.92±8.08	0.543
SFT(mm)	16.94±2.96	18.25±2.89	0.001*
MAMC(cm)	19.22±2.66	20.55±2.46	0.345
BMI(kg/m ²)	22.72±2.84	23.45±2.17	0.001*

*P<0.001 considered as significant

Figure.1 Age and sex wise distribution of study subjects




Figure.2 Mean values of biochemical parameters at 10% and 20% barley intervention





Fig. 3 Mean values of nutritional status parameters at 10% and 20% barley intervention



REFERENCE

1. Laura E Newton. Kidney Disease. In, Douglas C Hamburger, Jamey D (ed). Hand book of clinical Nutrition, 4th edition. New Delhi, Elsevier Publishers, 2009; 510-522.
2. Donna Zimmaro Bliss. Dietary fiber in conservative management of chronic renal disease. Pediatr Nephrol 2004; 19:1069-70.
3. Malan hot P. Natural health remedies and alternative medicine articles and information health benefits of cereals. Html | 4. Shakunthala Manay N. Shadakshraswamy. Barley. In. Shakunthala Manay N. Shadakshraswamy(ed). Foods Facts and principles, 1st edition. New Delhi, New Age International Publishers, 1997; 248-50.
5. McIntosh's GH. Whyte cholesterol in hypercholesterolemia men. Am J Clin Nutr.1991; 53:1205-9.
6. Madhavi reddy M, Raja reddy P, Raghavendra Prasad BN, Munilakshmi U. Grain and milling quality of barley and their suitability for preparation of traditional South Indian products. International Journal of Scientific Research, IOSR Journal of Pharmacy, Volume :4, Issue:2, February,2014, p: 23-27.
7. Scott D Cohen, Paul L Kimmel. Nutritional status, psychological status and survival in hemodialysis patients. Nutrition and kidney disease: A new Era, contrib.Nephrol.Basel, Karger, 2007; 5:1-17.
8. National Kidney Foundation: K/DOQI CLINICAL practice guidelines on hypertension and antihypertensive agents in chronic kidney disease.2004.from <http://www.kidney.org>>kidney disease.
9. National Kidney Foundation: K/DOQI CLINICAL practice guidelines and clinical practice recommendations for diabetes and chronic kidney disease.2007.<http://www.diabetes.ca/files/for-professionals/diet-kidney-disease>.
10. Mahdi GS,Abdal M, Behera BC, Verma N, Sonone A, Makhi U. barley is a health full food: a review. Electron J Environ Agric Food Chem 2008;&2686-94.
11. Lupton JR, Robinson MC, Morin JL.Cholesterol lowering effects of barley bran flour and oil. J Am diet Assoc1994; 94:65-70.
12. Anderson JW, Bidges SR. Hypercholesterolemia effects of oat bran in humans. In: Wood PJ, editor. Oat bran.St. Paul: American Association of cereal chemists; 1993.P.139-157.
13. Wilson TA,Nicolas RJ, delancy B, Chadwell, Moolchandani V, Kotyla T et al. reduced and high molecular weight barley B-glucan decrease plasma National kidney Foundation: K/DOQI Clinical practice guidelines for nutrition in National kidney failure. Am J Kidney Dis 2001; 37:s66-s70.
14. Pins J, Keenan JM, Curry LL, Goulson MJ, Kolberg LW. Extracted barley beta-glucan improves CVD risk factors and other biomarkers in a population of generally healthy hypercholesterolemic men and women.Prev Control 2005; 1:131.
15. Wood PJ. Relationships between soluble properties of cereal beta-glucans and physiological effects- a review. Trends food sci technol 2002; 13:313-20.
16. Jenkins AL, Jenkins DJA, Zdravkovic U, Wursch P, Vuksan V. Depression of the glycemic index by high levels of b-glucan fiber in two functional foods tested in type 2 diabetes. Eur J Clin Nutr.2002; 56:622-8.
17. Wursch P, Pi-Sunyer FX. The role of viscous

13/14