

Multiple Splenic Artery Aneurysms Secondary to Extra Hepatic Portal Vein Obstruction

Y.C. MENJUNATHA, K.N.V. PRASAD, Y.C. BEEREGOWDA, BHASKARAN A.

Key Words: Splenic artery, Aneurysm, Extra hepatic

INTRODUCTION

A 14 year old girl presented with vague abdominal pain, recurrent, well tolerated haematemesis and feeling of presence of a lump in the abdomen. Her abdominal examination revealed gross splenomegaly. The laboratory tests, haematological counts, liver enzymes, platelet counts and the international normalized ratio were within the normal limits. Upper gastrointestinal endoscopy showed grade II oesophageal varices. Ultrasonography showed that the portal vein was replaced by multiple, dilated, collateral venous channels around the porta, with collaterals around the pancreas and splenic hilum. There was splenomegaly with a suspicious aneurysm of the splenic artery at the hilum. She was recommended Computed Tomography (CT) angiography, as it was difficult to differentiate aneurysm from collaterals at the splenic hilum on ultrasonography. CT angiography revealed multiple aneurysms in the main splenic artery, at the splenic hilum and at intra parenchymal branches, with largest being seen at the splenic hilum, which measured 18 x 15mm [Table/Fig-1A and B].



Portovenous phase CT showed multiple dilated venous collaterals replacing the main portal vein with multiple collaterals around the pancreatic head and the splenic hilum. There was gross splenomegaly [Table/Fig-1C and D].

The aneurysms were indistinguishable from the dilated collaterals

on portovenous phase CT. A diagnosis of extra hepatic portal vein obstruction with splenomegaly and multiple splenic aneurysms was made. Splenectomy, aneurysmectomy and a splenorenal shunt were performed. The post operative period was uneventful.

DISCUSSION

Splenic Artery Aneurysms (SAAs) occur in around 4% of the patients with chronic liver disease and more commonly in females. The risk factors for the SAAs are arterial fibrodysplasia, multiparity, arteriosclerosis, pancreatitis, trauma, chronic liver disease and portal hypertension, the latter two being the commonest, as was in our case. The most common location of the aneurysm was the distal third of the splenic artery and the intra parenchymal branches. The increased concentration of few hormones may play a major role in the development of the aneurysms. The risk of the SAAs may increase with an increase in the patients' age, the spleen size and the portal vein pathologies. The clinical significance of the SAAs is prevention of the fatal complications of the rupture, although they are rare. It is important to diagnose SAAs before liver transplantation because of their propensity for rupture in the post transplant period. Identifying the SAAs in the setting of the large dilated collaterals in the splenic hilum can be difficult by ultrasonography and by portovenous phase CT scan [1]. A dedicated arterial and venous phase CT should be performed to



[Table/Fig-1A]: CT angiography axial section in 14 year old girl showing tortuous splenic artery with large aneurysm in the distal part of the splenic artery (arrow) and another smaller one in the intra parenchymal branch (arrow head).

[Table/Fig-1B]: Coronal, reformat CT angiography image showing splenic aneurysm.

THE UNIVERSITY OF CHICAGO DEPARTMENT OF CHEMISTRY FACULTY OF SCIENCE

THE UNIVERSITY OF CHICAGO
DEPARTMENT OF CHEMISTRY
FACULTY OF SCIENCE

THE UNIVERSITY OF CHICAGO
DEPARTMENT OF CHEMISTRY
FACULTY OF SCIENCE

THE UNIVERSITY OF CHICAGO
DEPARTMENT OF CHEMISTRY
FACULTY OF SCIENCE

THE UNIVERSITY OF CHICAGO
DEPARTMENT OF CHEMISTRY
FACULTY OF SCIENCE

THE UNIVERSITY OF CHICAGO
DEPARTMENT OF CHEMISTRY
FACULTY OF SCIENCE

THE UNIVERSITY OF CHICAGO
DEPARTMENT OF CHEMISTRY
FACULTY OF SCIENCE

THE UNIVERSITY OF CHICAGO
DEPARTMENT OF CHEMISTRY
FACULTY OF SCIENCE

THE UNIVERSITY OF CHICAGO
DEPARTMENT OF CHEMISTRY
FACULTY OF SCIENCE

THE UNIVERSITY OF CHICAGO
DEPARTMENT OF CHEMISTRY
FACULTY OF SCIENCE

THE UNIVERSITY OF CHICAGO
DEPARTMENT OF CHEMISTRY
FACULTY OF SCIENCE