

## Original Research Article

# Neutrophil to lymphocyte ratio in diagnosing acute cholecystitis: a retrospective cohort study in a tertiary rural hospital

Karthik Hareen T. V. K.\*, Bhaskaran A., Jaswanthi A. R.

Department of Surgery, SDU Medical College, Kolar, Karnataka, India

**Received:** 10 October 2016

**Accepted:** 09 November 2016

### \*Correspondence:

Dr. Karthik Hareen T. V. K.,

E-mail: [karthikhareen08@gmail.com](mailto:karthikhareen08@gmail.com)

**Copyright:** © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## ABSTRACT

**Background:** Acute cholecystitis accounts for most of the hospital admissions related to gastrointestinal diseases. In approximately 90% of patients, inflammation develops due to obstruction of the cystic duct by one or more gall-tones. Delayed management can lead to increased morbidity, due to progression to severe cholecystitis, such as gangrenous change, abscess formation, and gallbladder perforation. The neutrophil to lymphocyte ratio is derived from the counts of circulating neutrophils and lymphocytes, both of which are major leukocyte subpopulations. The inflammation triggered release of arachidonic acid metabolites and platelet-activating factors results in neutrophilia, and cortisol induced stress results in relative lymphopenia, and thus, the neutrophil to lymphocyte ratio accurately represents the underlying inflammatory process. The objective of this study was to evaluate the utility of the neutrophil to lymphocyte ratio (NLR) as a prognostic indicator in patients with cholecystitis. And to identify a relevant NLR value that discriminates between simple and severe cholecystitis.

**Methods:** A retrospective analysis of patient data compiled by assessment of operative findings documented and patient follow up recorded over the period encompassing three years (2015-2012) at RL Jalappa Hospital and Research Centre, Tamaka, Kolar, Karnataka, India.

**Results:** Our study population comprised of 250 patients with simple cholecystitis (69.63%) and 109 patients with severe cholecystitis (30.36%). The NLR of 3.0 could predict severe cholecystitis with 70.5% sensitivity and 70.0% specificity. A higher NLR ( $\geq 3.0$ ) was significantly associated with older age ( $p = 0.001$ ), male gender ( $p = 0.001$ ), admission via the emergency department ( $p < 0.001$ ), longer operation time ( $p < 0.001$ ), higher incidence of postoperative complications ( $p = 0.056$ ), and prolonged length of hospital stay (LOS) ( $p < 0.001$ ).

**Conclusions:** NLR  $\geq 3.0$  was significantly associated with severe cholecystitis and prolonged LOS in patients undergoing cholecystectomy. Therefore, preoperative NLR in patients undergoing cholecystitis due to cholecystitis seemed to be a useful surrogate marker for severe cholecystitis.

**Keywords:** Cholecystitis, Cholecystectomy, NLR, Severity

## INTRODUCTION

This is an exciting time in medicine. The pace of Acute cholecystitis is one of the most common conditions requiring an emergency care.<sup>1</sup> In majority of patients, inflammation develops due to obstruction of the cystic duct by gallstones.<sup>2</sup> Delayed management can worsen the

condition leading to increased morbidity, due to progression to severe cholecystitis, such as gangrenous changes, abscess formation, gallbladder perforation, etc.<sup>3,4</sup>

Diagnosis in patients with severe cholecystitis is often tough to establish both clinically and radio logically,

since the clinical manifestations can vary, and imaging studies are often equivocal.<sup>5</sup> However, marked contrasts in the morbidity and mortality rates have been observed between patients with simple cholecystitis and severe cholecystitis.<sup>3,6</sup> Therefore, accurate detection and proper management of patients who are at risk of developing severe cholecystitis is very much essential in preventing associated complications.

To predict the prognosis of inflammatory conditions, several scores have been suggested, including the neutrophil to lymphocyte ratio (NLR), modified Glasgow prognostic score, platelet to lymphocyte ratio, etc.<sup>7,8</sup> Of these, the NLR received greater importance, since it is simple to calculate, and involves no additional cost, as it uses results from a standard complete blood counts. The NLR is derived from the ratios of counts of circulating neutrophils and lymphocytes, both of which are major leukocyte subpopulations. Inflammation usually triggers the release of arachidonic acid metabolites and platelet-activating factors which results in neutrophilia. Simultaneously cortisol-induced stress results in relative lymphopenia, and thus, the NLR accurately represents the ongoing inflammatory process.<sup>9</sup>

The objective of this study was to evaluate the utility of the NLR as a prognostic indicator in patients with cholecystitis, and to identify a relevant ratio value that discriminates between simple and severe cholecystitis.

## METHODS

A retrospective analysis of patient data compiled by assessment of operative findings documented and patient follow up recorded over the period encompassing three years (2015-2011) at RL Jalappa hospital and research centre, Tamaka, Kolar. During the study period, 399 cholecystectomies were performed, majority of which were done through laparoscopic approach. To identify gallbladder inflammation, we only selected 371 patients in whom the pathology reports indicated the presence of cholecystitis. Finally, a total of 359 patients were selected for inclusion in this study. Each patient's NLR was calculated at admission time.

### Terminology and definitions

Cholecystitis is usually defined by a histological finding of inflammatory infiltrate of the gallbladder wall. Severe cholecystitis is defined as cholecystitis complicated by secondary changes, which include haemorrhage, gangrenous changes, emphysema, or perforation, etc and/or when the pathological examination indicated xantho-granulomatous cholecystitis.


### Statistical analysis

Data were described as means  $\pm$  standard deviations, or as medians and ranges. Continuous variables were compared using the independent t-test, while categorical

variables were compared using the chi-squared test. Multiple regression analyses were performed using a proportional hazards model to identify factors independently associated with the LOS greater than the 80th percentile after cholecystectomy, and to estimate corresponding odds ratio (OR) in 95% confidence intervals (CI). Statistical analysis was performed using SPSS version 15.0 (SPSS Inc., Chicago, IL). Statistical significance was accepted for P-values less than 0.05.


## RESULTS

A total of 359 patients who underwent cholecystectomy owing to symptoms of cholecystitis during the study period were included. The median age was 55.2 (13 - 91) years, and the patients comprised 165 women (45.96%) and 194 men (54.03%).



**Figure 1: Sex ratio.**


Of these, 294 patients (81.89%) exhibited calculous cholecystitis and 65 patients (18.1%) exhibited acalculous cholecystitis.



**Figure 2: Presence of calculus.**

Pathologic examination confirmed the presence of simple cholecystitis in 298 patients (83%) and severe cholecystitis in 61 patients (16.99%). Regarding the admission, 283 patients (78.83%) were admitted via the emergency department (ED) and 76 (21.16%) via the outpatient clinic.

An ROC curve was established to determine the cut-off value for preoperative NLR, which could discriminate between simple and severe cholecystitis.



**Figure 3: Mode of admission.**

The ROC area under the curve was 0.775. With an NLR value of 3.090, the sensitivity and specificity values were 70.5% and 70.0%, respectively. Therefore, we considered 3.0 as the cutoff value, and divided the patient population into two groups: those with preoperative NLR values less than 3.0 ( $n = 216$ ) and those with NLR values equal to or greater than 3.0 ( $n = 143$ ).

When comparing preoperative variables, the two groups showed significant differences in age, sex, severity of cholecystitis, serum leukocyte count, presenting symptoms and admission route. The higher NLR ( $NLR \geq 3$ ) group included more patients who had advanced age ( $p = 0.001$ ), number of males ( $p = 0.001$ ), with severe cholecystitis ( $p < 0.001$ ) and with higher leukocyte count ( $p < 0.001$ ).

## DISCUSSION

Severe cholecystitis is associated with more adverse clinical features than simple cholecystitis. So, prompt detection of the severe cholecystitis and intervention before its further advancement is essential to avoid complications. These patients are at an increased risk of injury to the biliary ducts, injury to the right hepatic artery during surgery, etc.<sup>10</sup>

In the current study, we elicited that a preoperative NLR of 3.0 has the usefulness to differentiate between simple and severe cholecystitis. Therefore, in patients whom the CT findings are ambiguous and are at a risk of progressing from simple to severe cholecystitis, NLR could be used to determine surgical priority.

Patients with inflammatory and other associated malignant conditions usually present with elevated NLR as a manifestation of the systemic response. The underlying molecular basis of NLR has found that there is an elevation in the levels of pro-inflammatory cytokines (e.g., IL-1ra, IL-6, IL-7, IL-8, IL-12) in the plasma of these patients.<sup>11-13</sup> Moreover, high peritumoral infiltration of macrophages was observed in cancer patients with elevated NLR.<sup>12</sup> Therefore, elevated NLR appears to be a good indicator of up-regulation of the innate immune response.

The representative forms of severe cholecystitis are gangrenous cholecystitis and gallbladder perforation. Gangrenous cholecystitis occurs in up to 30-35% of patients with cholecystitis.<sup>5</sup> This occurs because the inflammation causes interruption of blood flow to the gallbladder which results in gangrenous changes. The mortality rate was reported to be up to 22% in these patients, and it is directly related to other severe complications, such as gallbladder perforation, abscess formation, peritonitis, etc.<sup>14</sup>

This study shows the usefulness of preoperative NLR in predicting the prognosis and in determining the operative priority in patients with cholecystitis. Patients with severe cholecystitis have higher incidences of postoperative complications and a prolonged length of hospital stay (LOS).<sup>16-19</sup> Early cholecystectomy showed decrease LOS in patients with acute severe cholecystitis.<sup>20</sup>

Therefore, NLR helps in prioritizing the patients for surgery thus reducing the post-operative morbidity and LOS. Similarly, operation time was prolonged in the high NLR group than in the low NLR group.

Here we determined the cut-off value of severe cholecystitis as 3.0 based on the ROC curve analysis. This NLR value of 3.0 had an acceptable reliability in the analysis (70.6% sensitivity and a specificity of 70.1%). Therefore, we believe that the NLR cut-off value of 3.0 is suitable and consistent.<sup>27-30</sup> However, further study is required to validate the cut-off value and to more precisely identify an optimal NLR with the perfect and suitable prognostic power in cholecystitis.

Limitations of this study were most of those common to all database research. This study involves a retrospective review of prospectively collective data. Thus, these results should be confirmed by prospective trials. Moreover, patient distribution was not well-balanced according to the NLR cut-off value. In addition, the incidence of acalculous cholecystitis herein is more (18.1%) which did not fall within the general range of 2% to 15% as noted by other studies. Acute acalculous cholecystitis has generally shown poor prognosis than acute calculous cholecystitis.

## CONCLUSION

Routine preoperative NLR calculation in patients with cholecystitis not only provided a simple means of identifying patients with severe cholecystitis, but also served as a surrogate marker for predicting prolonged LOS. It was found that the patients with cholecystitis could be divided into low risk group for severe cholecystitis ( $NLR < 3.0$ ) and high risk group for severe cholecystitis ( $NLR \geq 3.0$ ) according to the NLR value at the time of admission. This kind of approach of determining the operative priority based on the NLR value is expected to induce a good surgical outcome by

prioritizing the sickest to be treated first principle and enabling proper perioperative management.

*Funding: No funding sources*

*Conflict of interest: None declared*

*Ethical approval: The study was approved by the institutional ethics committee*

## REFERENCES

1. Russo MW, Wei JT, Thiny MT, Gangarosa LM, Brown A, Ringel Y, et al. Digestive and liver diseases statistics. *Gastroenterology*. 2004;126(5):1448-53.
2. Charalel RA, Jeffrey RB, Shin LK. Complicated cholecystitis: the complementary roles of sonography and computed tomography. *Ultrasound Q*. 2011;27(3):161-70.
3. Ahmad MM, Macon WL. Gangrene of the gallbladder. *Am Surg*. 1983;49(3):155-8.
4. Wilson AK, Kozol RA, Salwen WA, Manov LJ, Tennenberg SD. Gangrenous cholecystitis in an urban VA hospital. *J Surg Res*. 1994;56(5):402-4.
5. Bennett GL, Rusinek H, Lisi V, Israel GM, Krinsky GA, Slywotzky CM, et al. CT findings in acute gangrenous cholecystitis. *AJR Am J Roentgenol*. 2002;178(2):275-81.
6. Morfin E, Ponka JL, Brush BE. Gangrenous cholecystitis. *Arch Surg*. 1968;96(4):567-73.
7. Kwon HC, Kim SH, Oh SY, Lee S, Lee JH, Choi HJ, et al. Clinical significance of preoperative neutrophil-lymphocyte versus platelet-lymphocyte ratio in patients with operable colorectal cancer. *Biomarkers: BiochemIndicat Expo Response Susceptibility Chem*. 2012;17(3):216-22.
8. Proctor MJ, Morrison DS, Talwar D, Balmer SM, Fletcher CD, O'Reilly DS, et al. A comparison of inflammation-based prognostic scores in patients with cancer. A glasgow inflammation outcome study. *Eur J Cancer*. 2011;47(17):2633-41.
9. Tamhane UU, Aneja S, Montgomery D, Rogers EK, Eagle KA, Gurm HS. Association between admission neutrophil to lymphocyte ratio and outcomes in patients with acute coronary syndrome. *Am J Cardiol*. 2008;102(6):653-7.
10. Davis B, Castaneda G, Lopez J. Subtotal cholecystectomy versus total cholecystectomy in complicated cholecystitis. *Am Surg*. 2012;78(7):814-7.
11. Kantola T, Klintrup K, Vayrynen JP, Vornanen J, Bloigu R, Karhu T, et al. Stage dependent alterations of the serum cytokine pattern in colorectal carcinoma. *Br J Cancer*. 2012;107(10):1729-36.
12. Motomura T, Shirabe K, Mano Y, Muto J, Toshima T, Umemoto Y, et al. Neutrophil-lymphocyte ratio reflects hepatocellular carcinoma recurrence after liver transplantation via inflammatory microenvironment. *J Hepatol*. 2013;58(1):58-64.
13. Ishizuka M, Nagata H, Takagi K, Iwasaki Y, Kubota K. Combination of platelet count and neutrophil to lymphocyte ratio is a useful predictor of postoperative survival in patients with colorectal cancer. *Br J Cancer*. 2013;109(2):401-7.
14. Grant RL, Tie ML. False negative biliary scintigraphy in gangrenous cholecystitis. *Australas Radiol*. 2002;46(1):73-5.
15. Derici H, Kara C, Bozdag AD, Nazli O, Tansug T, Akca E. Diagnosis and treatment of gallbladder perforation. *World J Gastroenterol WJG*. 2006;12(48):7832-6.
16. Bingener J, Stefanidis D, Richards ML, Schwesinger WH, Sirinek KR. Early conversion for gangrenous cholecystitis: impact on outcome. *Surg Endosc*. 2005;19(8):1139-41.
17. Choi SB, Han HJ, Kim CY, Kim WB, Song TJ, Suh SO, et al. Early laparoscopic cholecystectomy is the appropriate management for acute gangrenous cholecystitis. *Am Surg*. 2011;77(4):401-6.
18. Falor AE, Zobel M, Kaji A, Neville A, De Virgilio C. Admission variables predictive of gangrenous cholecystitis. *Am Surg*. 2012;78(10):1075-8.
19. Tsushimi T, Matsui N, Takemoto Y, Kurazumi H, Oka K, Seyama A, et al. Early laparoscopic cholecystectomy for acute gangrenous cholecystitis. *Surg Laparosc Endosc Percutan Tech*. 2007;17(1):14-8.
20. Murata A, Okamoto K, Matsuda S, Kuwabara K, Ichimiya Y, Matsuda Y, et al. Multivariate analysis of factors influencing length of hospitalization and medical costs of cholecystectomy for acute cholecystitis in Japan: a national database analysis. *Keio J Med*. 2013;62(3):83-94.
21. Guthrie GJ, Charles KA, Roxburgh CS, Horgan PG, McMillan DC, Clarke SJ. The systemic inflammation-based neutrophil-lymphocyte ratio: experience in patients with cancer. *Crit Rev Oncol Hematol*. 2013;88(1):218-30.
22. Carruthers R, Tho LM, Brown J, Kakumanu S, McCartney E, McDonald AC. Systemic inflammatory response is a predictor of outcome in patients undergoing preoperative chemoradiation for locally advanced rectal cancer. *Colorectal Dis Offic J Assoc Coloproctology Great Brit Ireland*. 2012;14(10):e701-7.
23. Cedres S, Torrejon D, Martinez A, Martinez P, Navarro A, Zamora E, et al. Neutrophil to lymphocyte ratio (NLR) as an indicator of poor prognosis in stage IV non-small cell lung cancer. *Clin Translational Oncol: Offic Publ Fed Spanish Oncol Soc Nat Canc Inst Mexico*. 2012;14(11):864-9.
24. Chua TC, Chong CH, Liauw W, Zhao J, Morris DL. Inflammatory markers in blood and serum tumor markers predict survival in patients with epithelial appendiceal neoplasms undergoing surgical cytoreduction and intraperitoneal chemotherapy. *Ann Surg*. 2012;256(2):342-9.

25. Kishi Y, Kopetz S, Chun YS, Palavecino M, Abdalla EK, Vauthey JN. Blood neutrophil-to-lymphocyte ratio predicts survival in patients with colorectal liver metastases treated with systemic chemotherapy. *Ann Surg Oncol.* 2009;16(3):614-22.
26. Sharaiha RZ, Halazun KJ, Mirza F, Port JL, Lee PC, Neugut AI, et al. Elevated preoperative neutrophil: lymphocyte ratio as a predictor of postoperative disease recurrence in esophageal cancer. *Ann Surg Oncol.* 2011;18(12):3362-9.
27. Chiang SF, Hung HY, Tang R, Changchien CR, Chen JS, You YT, et al. Can neutrophil to lymphocyte ratio predict the survival of colorectal cancer patients who have received curative surgery electively? *Int J Colorectal Dis.* 2012;27(10):1347-57.
28. Malietzis G, Giacometti M, Askari A, Nachiappan S, Kennedy RH, Faiz OD, et al. A preoperative neutrophil to lymphocyte ratio of 3 predicts disease-free survival after curative elective colorectal cancer surgery. *Ann Surg.* 2013;260(2):287-92.
29. Jeong JH, Lim SM, Yun JY, Rhee GW, Lim JY, Cho JY, et al. Comparison of two inflammation-based prognostic scores in patients with unresectable advanced gastric cancer. *Oncology.* 2012;83(5):292-9.
30. Keizman D, Ish-Shalom M, Huang P, Eisenberger MA, Pili R, Hammers H, et al. The association of pre-treatment neutrophil to lymphocyte ratio with response rate, progression free survival and overall survival of patients treated with sunitinib for metastatic renal cell carcinoma. *Eur J Cancer.* 2012;48(2):202-8.

**Cite this article as:** Hareen TVK, Bhaskaran A, Jaswanthi AR. Neutrophil to lymphocyte ratio in diagnosing acute cholecystitis: a retrospective cohort study in a tertiary rural hospital. *Int Surg J* 2017;4:372-6.